

Mouse CD49d Monoclonal Antibody | anti-CD49d antibody

MOUSE ANTI HUMAN CD49d


CATALOG # AAA49454	HOST Mouse	APPLICATIONS Flow Cytometry, Functional Assay, Immunohistochemistry
------------------------------	----------------------	---

PRODUCT SYNONYMS


CD49d; Monoclonal Antibody; MOUSE ANTI HUMAN CD49d; anti-CD49d antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Number	44H6
Form/Format	Purified Purified IgG - liquid
Concentration	IgG concentration 1.0 mg/ml (varies by lot)
Sequence Length	1032
Applicable Applications for anti-CD49d antibody	FCM/FACS (Flow Cytometry), IHC (Immunohistochemistry)
Perservative Stabilisers	0.09% Sodium Azide Preparation
Histology Positive Control Tissue	Human Tonsil Buffer Solution
Target Species	Human
Preparation and Storage	Store at 4 degree C or at -20 degree C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 18 months from date of despatch.




Application Data
Staining of human peripheral blood lymphocytes with Mouse anti Human CD49d



Application Data
Staining of human peripheral blood lymphocytes with Mouse anti Human CD49d: FITC



Application Data
Staining of human peripheral blood lymphocytes with Mouse anti Human CD49d: Low Endotoxin



Application Data
Staining of human peripheral blood lymphocytes with Mouse anti Human CD49d:RPE

ADDITIONAL INFORMATION

Related Product Information for anti-CD49d antibody

Mouse anti Human CD49d monoclonal antibody, clone 44H6 recognizes human CD49d also known as integrin alpha-4 or VLA-4 subunit alpha. CD49d is a ~150kDa single pass type 1 transmembrane glycoprotein with seven FG-GAP repeats, characteristic of alpha integrins, in its extracellular domain. CD49d can be proteolytically cleaved to yield fragments of 80 and 70kDa (Hemler et al. 1987). CD49d associates with either CD29 to form VLA-4 or with Integrin beta-7 to form The Peyer patches-specific homing receptor LPAM-1, involved in the lymphocyte migration and homing to gut-associated lymphoid tissue (Sackstein 2006) through its interaction with MadCam-1, preferentially expressed on Peyer's patch high endothelial venules and postcapillary venules in lamina propria (Briskin et al. 1997) Mouse anti human CD49d, clone 44H6 is reported to induce homotypic aggregation of pre-B cell lines including HOON and NALM-6 (Letarte et al. 1993).

NCBI AND UNIPROT INFORMATION

NCBI GI #	67191027
NCBI GeneID	3676
NCBI Accession #	NP_000876.3
NCBI GenBank Nucleotide #	NM_000885.4
UniProt Accession #	P13612; Q7Z4L6; D3DPG4; AAB25486
Molecular Weight	21,536 Da
NCBI Official Full Name	integrin alpha-4

NCBI Official Synonym Full Names	integrin, alpha 4 (antigen CD49D, alpha 4 subunit of VLA-4 receptor)
NCBI Official Symbol	ITGA4
NCBI Official Synonym Symbols	IA4; CD49D
NCBI Protein Information	integrin alpha-4; 269C wild type; CD49 antigen-like family member D; VLA-4 subunit alpha; antigen CD49D, alpha-4 subunit of VLA-4 receptor; integrin alpha 4; integrin alpha-4 subunit; integrin alpha-IV; very late activation protein 4 receptor, alpha 4 subunit
UniProt Protein Name	Integrin alpha-4
UniProt Gene Name	ITGA4
UniProt Synonym Gene Names	CD49D
UniProt Entry Name	ITA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd49d-antibody-clone-44h6-flow-cytometry-immunohistochemistry-elisa/49454>