

Mouse CD18 Monoclonal Antibody | anti-CD18 antibody

MOUSE ANTI HUMAN CD18 (ACTIVATION EPI TOPE):RPE

CATALOG # AAA49487	HOST Mouse	APPLICATIONS Flow Cytometry, Functional Assay
------------------------------	----------------------	---


PRODUCT SYNONYMS

CD18; Monoclonal Antibody; MOUSE ANTI HUMAN CD18 (ACTIVATION EPI TOPE):RPE; anti-CD18 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Number	MEM-148
Form/Format	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilised
Sequence Length	769
Applicable Applications for anti-CD18 antibody	FCM/FACS (Flow Cytometry)
Reconstitution	Reconstitute with 1 ml distilled water Perservative Stabilisers
Buffer Solution	Phosphate buffered saline
Target Species	Human
Preparation and Storage	Prior to reconstitution store at 4 degree C. Following reconstitution store at 4 degree C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 12 months from date of reconstitution.

IMAGES




Application Data

Staining of human peripheral blood granulocytes with Mouse anti Human CD18 (Activation Epi tope):RPE




Application Data

Staining of human peripheral blood granulocytes with Mouse anti Human CD18 (ACT EPI TOPE



Application Data

Staining of human peripheral blood monocytes with Mouse anti Human CD86 (Activation Epitope): Azide free



Application Data

Staining of human peripheral blood monocytes with Mouse anti Human CD18 (Activation Epitope): Biotin

ADDITIONAL INFORMATION

Related Product Information for anti-CD18 antibody

Mouse anti Human CD18 (Activation Epitope) antibody, clone MEM-148 recognizes an epitope on the human CD18 molecule that is hidden in the CD11/CD18 heterodimer on resting cells. Clone MEM-148 binds very weakly to resting peripheral blood leukocytes and strongly to all leukocytes upon cellular activation. The epitope recognized by Mouse anti Human CD18 (Activation Epitope) antibody, clone MEM-148 is also exposed during dissociation of the CD11/CD18 by low pH.

NCBI AND UNIPROT INFORMATION

NCBI GI #	89191865
NCBI GeneID	3689
NCBI Accession #	NP_000202.2
NCBI GenBank Nucleotide #	NM_000211.3
UniProt Accession #	P05107; Q16418; Q53HS5; Q9UD72; B3KTS8; D3DSM1
Molecular Weight	84,782 Da
NCBI Official Full Name	integrin beta-2
NCBI Official Synonym Full Names	integrin, beta 2 (complement component 3 receptor 3 and 4 subunit)
NCBI Official Symbol	ITGB2
NCBI Official Synonym Symbols	LAD; CD18; MF17; MFI7; LCAMB; LFA-1; MAC-1
NCBI Protein Information	integrin beta-2; cell surface adhesion glycoprotein (LFA-1/CR3/P150,959 beta subunit precursor); complement receptor C3 beta-subunit; integrin beta chain, beta 2; leukocyte cell adhesion molecule CD18; leukocyte-associated antigens CD18/11A, CD18/11B, CD18/11C
UniProt Protein Name	Integrin beta-2
UniProt Gene Name	ITGB2
UniProt Synonym Gene Names	CD18; MFI7
UniProt Entry Name	ITB2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd18-antibody-clone-mem-148-flow-cytometry-elisa/49487>