

Mouse CD32 Monoclonal Antibody | anti-CD32 antibody

MOUSE ANTI HUMAN CD32:RPE


CATALOG # AAA49565	HOST Mouse	APPLICATIONS Flow Cytometry, Functional Assay
------------------------------	----------------------	---

PRODUCT SYNONYMS


CD32; Monoclonal Antibody; MOUSE ANTI HUMAN CD32:RPE; anti-CD32 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Number	AT10
Form/Format	Purified IgG conjugated to R. Phycoerythrin (RPE) - liquid
Sequence Length	317
Applicable Applications for anti-CD32 antibody	FCM/FACS (Flow Cytometry)
Perservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 5% Sucrose Preparation
Immunogen	K562 cell line Fusion Partners
Target Species	Human
Preparation and Storage	Store at 4 degree C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 12 months from date of despatch.




Application Data
Staining of human peripheral blood granulocytes with Mouse anti Human CD32:Alexa Fluor 647




Application Data
Staining of human peripheral blood granulocytes with Mouse anti Human CD32



Application Data
Staining of human peripheral blood granulocytes with Mouse anti Human CD32:RPE



Application Data
Staining of human peripheral blood granulocytes with Mouse anti Human CD32:FITC



Application Data
Staining of human peripheral blood monocytes with Mouse anti Human CD32: Biotin

ADDITIONAL INFORMATION

Related Product Information for anti-CD32 antibody

Mouse anti Human CD32 antibody, clone AT10 recognizes the human CD32 antigen, a 40kDa glycoprotein that acts as a low affinity receptor for IgG (also known as Fc gamma RII). The antigen mediates several functions including endocytosis, activation of secretion, cytotoxicity and immunomodulation. CD32 is expressed by B cells, monocytes, granulocytes and platelets. Clone AT10 blocks the binding of IgG to Fc gamma RII (Larsson et al. 1997).

NCBI AND UNIPROT INFORMATION

NCBI GI #	210031822
NCBI GeneID	2212
NCBI Accession #	NP_001129691.1
NCBI GenBank Nucleotide #	NM_001136219.1
Molecular Weight	34,930 Da
NCBI Official Full Name	low affinity immunoglobulin gamma Fc region receptor II-a isoform 1
NCBI Official Synonym Full Names	Fc fragment of IgG, low affinity IIa, receptor (CD32)
NCBI Official Symbol	FCGR2A
NCBI Official Synonym Symbols	CD32; FCG2; FcGR; CD32A; CDw32; FCGR2; IGFR2; FCGR2A1
NCBI Protein Information	low affinity immunoglobulin gamma Fc region receptor II-a; Immunoglobulin G Fc receptor II; fc-gamma RII-a; fc-gamma-RIIa; fcRII-a; igG Fc receptor II-a
UniProt Protein Name	Low affinity immunoglobulin gamma Fc region receptor II-a
UniProt Gene Name	FCGR2A
UniProt Synonym Gene Names	CD32; FCG2; FCGR2A1; IGFR2; IgG Fc receptor II-a; Fc-gamma-RIIa; FcRII-a
UniProt Entry Name	FCG2A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd32-antibody-clone-at10-flow-cytometry-elisa/49565>