

Mouse CD138 Monoclonal Antibody | anti-CD138 antibody

MOUSE ANTI HUMAN CD138:FITC

CATALOG # AAA49679	HOST Mouse	APPLICATIONS Flow Cytometry, Functional Assay
------------------------------	----------------------	---


PRODUCT SYNONYMS

CD138; Monoclonal Antibody; MOUSE ANTI HUMAN CD138:FITC; SYNDECAN-1; SDC; anti-CD138 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Number	B-A38
Form/Format	FITC Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid
Concentration	IgG concentration 0.1 mg/ml (varies by lot)
Sequence Length	310
Applicable Applications for anti-CD138 antibody	FCM/FACS (Flow Cytometry)
Perservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Preparation	Purified IgG prepared by ion exchange chromatography
Buffer Solution	Phosphate buffered saline
Target Species	Human
Immunogen	U266 cell line
Fusion Partners	Spleen cells from immunised BALB/c (Ifba Credo) mice were fused with cells of the mouse X63/Ag.8653 myeloma cell line
Preparation and Storage	Store at 4 degree C or at -20 degree C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 18 months from date of despatch.

IMAGES



Application Data
Staining of U266 cell line with Mouse anti Human CD138: FITC



Application Data
Staining of U266 cell line with Mouse anti Human CD138: Alexa Fluor 647

ADDITIONAL INFORMATION

Related Product Information for anti-CD138 antibody	Mouse anti human CD138 antibody, clone B-A38 recognizes human CD138, also known as Syndecan-1 (SDC-1). CD138 is a member of the transmembrane heparan sulfate proteoglycan family (O'Connell et al. 2004, Sanderson et al. 2008). It is composed of a core protein (comprising 3 domains; a short cytoplasmic domain, a transmembrane domain, and a long extracellular domain) and covalently attached heparan sulfate chains (Sanderson et al. 2008). Syndecan-1 is expressed on the surface of plasma cells within the hematopoietic system and on the surface of mature epithelial cells (O'Connell et al. 2004). It acts as an extracellular matrix receptor, involved in many cellular functions, including cell binding, cell signalling and cytoskeletal organization through cell-cell adhesion and cell-matrix adhesion (Sanderson et al. 2008).
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	55749480
NCBI GeneID	6382
NCBI Accession #	NP_001006947.1
NCBI GenBank Nucleotide #	NM_001006946.1
UniProt Accession #	P18827; Q53QV0; Q546D3; Q96HB7; D6W523; AAB21434; AAB23935
Molecular Weight	32,462 Da
NCBI Official Full Name	syndecan-1
NCBI Official Synonym Full Names	syndecan 1
NCBI Official Symbol	SDC1
NCBI Official Synonym Symbols	SDC; CD138; SYND1; syndecan
NCBI Protein Information	syndecan-1; CD138 antigen; heparan sulfate proteoglycan fibroblast growth factor receptor; syndecan proteoglycan 1
UniProt Protein Name	Syndecan-1
UniProt Gene Name	SDC1
UniProt Synonym Gene Names	SDC; SYND1
UniProt Entry Name	SDC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

