

Goat CCR10 Polyclonal Antibody | anti-CCR10 antibody

GOAT ANTI HUMAN CCR10

CATALOG # AAA50063	HOST Goat	APPLICATIONS Western Blot, Immunohistochemistry, Flow Cytometry, Functional Assay, ELISA
------------------------------	---------------------	--


PRODUCT SYNONYMS

CCR10; Polyclonal Antibody; GOAT ANTI HUMAN CCR10; anti-CCR10 antibody

PRODUCT SPECIFICATIONS


Host	Goat
Clonality	Polyclonal
Isotype	IgG
Form/Format	Purified Purified IgG - liquid
Concentration	IgG concentration 1.0mg/ml (varies by lot)
Sequence Length	362
Applicable Applications for anti-CCR10 antibody	WB (Western Blot), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry), ELISA
Perservative Stabilisers	0.1% Sodium Azide 0.1% Bovine Serum Albumin
Histology Positive Control Tissue	Spleen or tonsil Buffer Solution
Antiserum Preparation	Antiserum to human CCR10 was raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared by affinity chromatography.
Target Species	Human
Preparation and Storage	Store at 4 degree C or at -20 degree C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 18 months from date of despatch.

IMAGES



Application Data

Paraffin embedded human tonsil stained with Goat anti Human CCR10 (AAA50063)



Application Data

Paraffin embedded human spleen stained with Goat anti Human CCR10 (AAA50063)

ADDITIONAL INFORMATION

Related Product Information for anti-CCR10 antibody	Goat anti Human CCR10 antibody recognizes human CCR10 (C-C chemokine receptor type 10), otherwise known as GPR2, (G-protein coupled receptor 2), a transmembrane glycoprotein and member of the G-protein coupled receptor 1 family, expressed by melanocytes, dermal fibroblasts, and dermal microvascular endothelial cells, which acts as a receptor for the skin-associated chemokine CCL27 (SCYA27/CTACK), and is implicated in T cell-mediated skin inflammation and possibly skin homeostasis.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	156104886
NCBI GeneID	2826
NCBI Accession #	NP_057686.2
NCBI GenBank Nucleotide #	NM_016602.2
UniProt Accession #	P46092; Q4V749; Q6T7X2; Q9NZG2
Molecular Weight	38,416 Da
NCBI Official Full Name	C-C chemokine receptor type 10
NCBI Official Synonym Full Names	chemokine (C-C motif) receptor 10
NCBI Official Symbol	CCR10
NCBI Official Synonym Symbols	GPR2
NCBI Protein Information	C-C chemokine receptor type 10; C-C CKR-10; CC chemokine receptor 10; CC-CKR-10; CCR-10; G protein-coupled receptor 2; G-protein coupled receptor 2
UniProt Protein Name	C-C chemokine receptor type 10
UniProt Gene Name	CCR10
UniProt Synonym Gene Names	GPR2; C-C CKR-10; CC-CKR-10; CCR-10
UniProt Entry Name	CCR10_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ccr10-antibody-western-blot-immunohistochemistry-flow-cytometry-elisa/50063>