

Mouse CD16 Monoclonal Antibody | anti-CD16 antibody

MOUSE ANTI HUMAN CD16:FITC

CATALOG # AAA50219	HOST Mouse	APPLICATIONS Flow Cytometry, Functional Assay
------------------------------	----------------------	---


PRODUCT SYNONYMS

CD16; Monoclonal Antibody; MOUSE ANTI HUMAN CD16:FITC; anti-CD16 antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Clone Number	KD1
Form/Format	FITC Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid
Concentration	IgG concentration 0.1mg/ml (varies by lot)
Sequence Length	233
Applicable Applications for anti-CD16 antibody	FCM/FACS (Flow Cytometry)
Perservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin Preparation
Buffer Solution	Phosphate buffered saline
Target Species	Human
Preparation and Storage	Store at 4 degree C or at -20 degree C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf Life: 18 months from date of despatch.

IMAGES



Application Data

Human peripheral blood lymphocytes stained with Mouse anti Human CD16: FITC(AAA50219)



Application Data

Human peripheral blood lymphocytes stained with Mouse anti Human CD16 followed by Rabbit F(ab')₂ anti Mouse IgG:FITC

ADDITIONAL INFORMATION

Related Product Information for anti-CD16 antibody	Mouse anti Human CD16 antibody, clone KD1 recognizes human CD16, a 50-65kD cell surface molecule, which is the low affinity receptor for IgG (FcR III). CD16 exists as a transmembranous form (Fc gammaRIIIA, or CD16A) and a glycosyl phosphatidylinositol (GPI) anchored form (Fc gammaRIIIB, or CD16B). CD16A is expressed by NK cells, some T cells, and macrophages (Moretta et al. 1990), whereas CD16B is primarily expressed by granulocytes (Bonecchi et al. 1999). Clone KD1 recognizes both forms of CD16 and will therefore recognize all cell types expressing CD16. Mouse anti Human CD16, clone KD1 can be used to identify CD16 in a range of species including bovine (Boysen et al. 2010) and ovine (Elh mouzi-Younes et al. 2010).
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	132814489
NCBI GeneID	2215
NCBI Accession #	NP_000561.3
NCBI GenBank Nucleotide #	NM_000570.4
Molecular Weight	26,216 Da
NCBI Official Full Name	low affinity immunoglobulin gamma Fc region receptor III-B isoform 2
NCBI Official Synonym Full Names	Fc fragment of IgG, low affinity IIIb, receptor (CD16b)
NCBI Official Symbol	FCGR3B
NCBI Official Synonym Symbols	CD16; FCG3; CD16b; FCGR3; FCR-10; FCRIII; FCRIIIb
NCBI Protein Information	low affinity immunoglobulin gamma Fc region receptor III-B; Fc-gamma receptor IIIb (CD 16); fc-gamma RIII-beta; fc-gamma RIIIb; igG Fc receptor III-1
UniProt Protein Name	Low affinity immunoglobulin gamma Fc region receptor III-B
UniProt Gene Name	FCGR3B
UniProt Synonym Gene Names	CD16B; FCG3; FCGR3; IGFR3; Fc-gamma RIII; Fc-gamma RIIIb; FcRIII; FcRIIIb
UniProt Entry Name	FCG3B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.