

Rabbit anti-Human Histone H3 Polyclonal Antibody | anti-HIST1H3F antibody

Histone H3 (Ac27) antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA50456	Rabbit	Human	Western Blot, Immunofluorescence, Chromatin Immunoprecipitation, Immunoprecipitation


PRODUCT SYNONYMS

Histone H3; Polyclonal Antibody; Histone H3 (Ac27) antibody; RABBIT ANTI HUMAN HISTONE H3 (Ac27); anti-HIST1H3F antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antiserum to human histone H3 (Ac27) was raised by repeated immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography on Protein A.
Form/Format	Purified IgG - liquid Phosphate buffered saline Antiserum to human histone H3 (Ac27) was raised by repeated immunization of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography on Protein A.
Applicable Applications for anti-HIST1H3F antibody	WB (Western Blot), IF (Immunofluorescence), ChIP (Chromatin immunoprecipitation)
Preparation and Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20 degree C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8 degree C for short term use (up to 4 weeks) and store the remaining aliquots at -20 degree C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use.

IMAGES



WB (Western Blot)

Western blot analysis of untreated Raji cell nuclear extract (20 ug; lane 1) or Raji cell nuclear extract treated with sodium butyrate (20 ug; lane 2) probed with histone H3 (Ac27) antibody (AAA50456) at a 1 ug/ul dilution.



IF (Immunofluorescence)

Immunofluorescence staining of HEK293 cells probed with histone H3 (Ac27) antibody (AAA50456) at a 1 ug/ml dilution (top) and DAPI (bottom).

ADDITIONAL INFORMATION

Related Product Information for anti-HIST1H3F antibody	Rabbit anti Human histone H3 (Ac27) antibody recognizes histone H3 when acetylated at lysine 27. Histone H3 is one of the four core histones that make up the nucleosome core particle. Nucleosomes are the smallest subunit of chromatin and are made up of 146 bp of DNA wrapped around an octamer comprised of pairs of the four core histones (H2A, H2B, H3, and H4) (Smith, 1991). Lysine residues within the N-terminal tail protruding from the histone core of the nucleosome are acetylated and deacetylated as part of gene regulation. Hyperacetylation of histone tails makes chromatin more accessible to DNA-binding proteins by weakening histone-DNA and nucleosome-nucleosome interactions (Kuo and Allis, 1998). Wide species cross-reactivity is expected from Rabbit anti Human histone H3 (Ac27) antibody based on sequence.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504281
NCBI GeneID	8350
NCBI Accession #	NP_003520.1
NCBI GenBank Nucleotide #	NM_003529.2
UniProt Accession #	P68431; P02295; P02296; P16106; Q6ISV8; Q6NWP8; Q6NWP9; Q6NXU4; Q71DJ3; Q93081; A0PJT7; A5PLR1
Molecular Weight	15,404 Da
NCBI Official Full Name	histone H3.1
NCBI Official Synonym Full Names	histone cluster 1, H3a
NCBI Official Symbol	HIST1H3A
NCBI Official Synonym Symbols	H3/A; H3FA
NCBI Protein Information	histone H3.1; histone H3/a; histone 1, H3a; H3 histone family, member A
UniProt Protein Name	Histone H3.1
UniProt Gene Name	HIST1H3A
UniProt Synonym Gene Names	H3FA
UniProt Entry Name	H31_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/histone-h3-antibody-western-blot-immunofluorescence-chromatin-immunoprecipitation-elisa/50456>