

Goat anti-Human GDF15 Polyclonal Antibody | anti-GDF15 antibody

Goat Polyclonal to Human GDF15

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA50879	Goat	Human	ELISA, Western Blot, Immunohistochemistry

PRODUCT SYNONYMS


GDF15; Polyclonal Antibody; Goat Polyclonal to Human GDF15; Anti-GDF15 Antibody (C-Terminus) IHC-plus; MIC1; MIC-1; NAG-1; NRG-1; NSAID-regulated gene 1 protein; PLAB; NSAID-activated gene 1 protein; Placental TGF-beta; PTGFB; GDF-15; PTGF-beta; Human GDF15; anti-GDF15 antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Human
Clonality	Polyclonal
Specificity	Human GDF15.
Purity/Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Form/Format	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Concentration	0.5 mg/ml (varies by lot)
Sequence Positions	286-299
Applicable Applications for anti-GDF15 antibody	ELISA, WB (Western Blot), IHC (Immunohistochemistry)
Target Species	Human
Immunogen	GDF15 antibody was raised against peptide with sequence C-QKTDGTVSLQTYDD from the C-terminus of the protein sequence according to (NP_004855.2).
Family/Subfamily	TGF beta/not assigned-TGF beta.
Disclaimer	Due to the highly specific nature of antibodies and antigens, we cannot predict or be held responsible with respect to how this antibody will behave in your systems. Researchers using this antibody should conduct optimization studies to achieve the most optimal result possible for their intended application.


Recommended Immunohistochemistry Protocol	The following protocol is a recommendation only, and AAA Biotech makes no guarantee of the results: Tissue Preparation: Formalin fixation and embedding in paraffin wax. Tissue Sectioning: Make 4-um sections and place on pre-cleaned and charged microscope slides. Heat in a tissue-dryingoven for 45 minutes at 60°C. Deparaffinization: Wash dry slides in 3 changes of xylene - 5 minutes each @ RT Rehydration: Wash slides in 3 changes of 100% alcohol - 3 minutes each @ RT Wash slides in 2 changes of 95% alcohol - 3 minutes each @ RT Wash slides in 1 change of 80% alcohol - 3 minutes @ RT Rinse slides in gentle running distilled water - 5 minutes @ RT Antigen retrieval: Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes Remove from heat and let stand at room temperature in buffer - 20 minutes Rinse in 1X TBS with Tween (TBST) -1 minute @ RT Immunostaining: (Do not allow tissues to dry at any time during the staining procedure) Apply a universal protein block - 20 minutes @ RT Drain protein block from slides, apply diluted primary antibody - 45 minutes @ RT Rinse slides in 1 X TBST - 1 minute @ RT Apply a biotinylated secondary antibody appropriate for the primary antibody - 30 minutes @ RT Rinse slides in 1X TBST -1 minute @ RT Apply alkaline phosphatase streptavidin - 30 minutes @ RT Rinse slides in 1X TBST -1 minute @ RT Apply alkaline phosphatase chromogen substrate - 30 minutes @ RT Wash slides in distilled water - 1 minute @ RT Dehydrate: (This method should only be used if the chromogen substrate is alcohol insoluble (e.g. Vector Red, DAB) Wash slides in 2 changes of 80% alcohol - 1 minute each @ RT Wash slides in 2 changes of 95% alcohol - 1 minute each @ RT Wash slides in 3 changes of 100% alcohol - 1 minute each @ RT Wash slides in 3 changes of xylene - 1 minute each @ RT Apply coverslip
Preparation and Storage	Aliquot and store at -20 degree C. Avoid freezing and thawing.

IMAGES




WB (Western Blot)

Antibody (0.1 ug/ml) staining of Human Prostate lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



IHC (Immunohistochemistry)

Anti-GDF15 antibody IHC of human prostate. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml.



IHC (Immunohistochemistry)

Anti-GDF15 antibody IHC of human placenta. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml.

ADDITIONAL INFORMATION

Related Product Information for anti-GDF15 antibody	Growth differentiation factor 15 (GDF15) is a protein belonging to the transforming growth factor beta superfamily that has a role in regulating inflammatory and apoptotic pathways in injured tissues and during disease processes. GDF15 is also known as TGF-PL, MIC-1, PDF, PLAB, and PTGFB. GDF15 mRNA is most abundant in the liver, with lower levels seen in some other tissues.
Product Categories/Family for anti-GDF15 antibody	Family: TGF beta

NCBI AND UNIPROT INFORMATION

NCBI GI #	153792495
NCBI GeneID	9518
NCBI Accession #	NP_004855.2
NCBI GenBank Nucleotide #	NM_004864.2
UniProt Accession #	Q99988 ; O14629 ; P78360 ; Q9BWA0 ; Q9NRT0
Molecular Weight	34,140 Da
NCBI Official Full Name	growth/differentiation factor 15
NCBI Official Synonym Full Names	growth differentiation factor 15
NCBI Official Symbol	GDF15
NCBI Official Synonym Symbols	PDF; MIC1; PLAB; MIC-1; NAG-1; PTGFB; GDF-15

NCBI Protein Information	growth/differentiation factor 15; NRG-1; PTGF-beta; placental TGF-beta; NSAID-activated gene 1 protein; NSAID-regulated gene 1 protein; prostate differentiation factor; macrophage inhibitory cytokine 1; placental bone morphogenetic protein; NSAID (nonsteroidal anti-inflammatory drug)-activated protein 1
UniProt Protein Name	Growth/differentiation factor 15
UniProt Gene Name	GDF15
UniProt Synonym Gene Names	MIC1; PDF; PLAB; PTGFB; GDF-15; MIC-1; NAG-1; NRG-1
UniProt Entry Name	GDF15_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/gdf15-antibody-western-blot-immunohistochemistry-elisa/50879>