

Rabbit anti-Human UBC / Ubiquitin C Polyclonal Antibody | anti-UBC antibody

Rabbit Polyclonal to Human UBC / Ubiquitin C


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA50980	Rabbit	Human	Western Blot, Immunohistochemistry

PRODUCT SYNONYMS


UBC / Ubiquitin C; Polyclonal Antibody; Rabbit Polyclonal to Human UBC / Ubiquitin C; Anti-UBC / Ubiquitin C Antibody (N-Terminus) IHC-plus; UBC; HUBCEP52; L40; HMG20; RPL40; Ubiquitin; Polyubiquitin-C; CEP52; Ubiquitin C; Human UBC; anti-UBC antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Ammonium sulfate precipitation
Form/Format	PBS, 0.09% sodium azide
Sequence Length	685
Applicable Applications for anti-UBC antibody	WB (Western Blot), IHC (Immunohistochemistry)
Target Species	Human
Immunogen Description	KLH conjugated synthetic peptide selected from the N-terminal region of human Ubiquitin.
Immunogen Type	Synthetic peptide
Immunogen	UBC / Ubiquitin C antibody was raised against KLH conjugated synthetic peptide selected from the N-terminal region of human Ubiquitin.
Antigen Modification	N-Terminus
Preparation and Storage	Maintain refrigerated at 2-8 degree C for up to 6 months. For long term storage store at -20 degree C.




WB (Western Blot)
Western blot of HeLa cell lysate.




WB (Western Blot)
Western blot analysis of anti-Ubiquitin Pab in HeLa cell lysate. Ubiquitin (Arrow) was detected using purified Pab. Secondary HRP-anti-rabbit was used for signal visualization with chemiluminescence.



IHC (Immunohistochemisry)
Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with Ubiquitin antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



IHC (Immunohiostchemistry)
Formalin-fixed and paraffin-embedded human Kidney tissue reacted with Ubiquitin antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



IHC (Immunohistochemistry)
Anti-UBC / Ubiquitin C antibody IHC of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 10 ug/ml.

NCBI AND UNIPROT INFORMATION

NCBI GI #	601984520
NCBI GeneID	7316
NCBI Accession #	NP_066289.3
NCBI GenBank Nucleotide #	NM_021009.6

UniProt Accession #	P0CG48; P02248; P02249; P02250; P62988; Q29120; Q6LBL4; Q6LDU5; Q8WYN8; Q91887; Q91888; Q9BWD6
Molecular Weight	77,039 Da
NCBI Official Full Name	polyubiquitin-C
NCBI Official Synonym Full Names	ubiquitin C
NCBI Official Symbol	UBC
NCBI Official Synonym Symbols	HMG20
NCBI Protein Information	polyubiquitin-C
UniProt Protein Name	Polyubiquitin-C
UniProt Gene Name	UBC
UniProt Entry Name	UBC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ubc-ubiquitin-c-antibody-western-blot-immunohistochemistry-elisa/50980>