

Rabbit TCF19 / TCF-19 Polyclonal Antibody | anti-TCF19 antibody

Rabbit Polyclonal to Human TCF19 / TCF-19

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA51469	Rabbit	Gibbon, Chimpanzee, Gorilla, Human, Monkey	Western Blot, Immunohistochemistry



PRODUCT SYNONYMS

TCF19 / TCF-19; Polyclonal Antibody; Rabbit Polyclonal to Human TCF19 / TCF-19; Anti-TCF19 / TCF-19 Antibody (aa259-274) IHC-plus; TCF19; SC1(TCF19)-7; SC1; TCF-19; Transcription factor 19; Transcription factor SC1; SC1(TCF19)-4; SC1(TCF19)-6; SC1-1; Human TCF19; anti-TCF19 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Gibbon, Chimpanzee, Gorilla, Human, Monkey
Clonality	Polyclonal
Specificity	Human TCF19
Purity/Purification	Immunoaffinity Purified
Form/Format	PBS, 0.05% BSA, 0.05% sodium azide
Concentration	0.5 mg/ml (varies by lot)
Sequence Length	345
Applicable Applications for anti-TCF19 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Target Species	Human
Immunogen Description	A synthetic peptide corresponding to amino acids 259-274 (KAPLTPTGNRRGRPRK) of human TCF19/SC1 was used as immunogen, GenBank no Q9Y242.
Immunogen Type	Synthetic peptide
Immunogen	TCF19 / TCF-19 antibody was raised against a synthetic peptide corresponding to amino acids 259-274 (KAPLTPTGNRRGRPRK) of human TCF19/SC1 was used as immunogen, GenBank no Q9Y242. Percent identity by BLAST analysis: Human, Chimpanzee, Gorilla, Gibbon, Monkey (100%); Rabbit, Horse (88%); Marmoset, Bovine, Pig (81%).
Antigen Modification	aa259-274
Preparation and Storage	Store at 4 degree C for short term applications. For long term storage, aliquot and store at -20 degree C.

IMAGES



IHC (Immunohistochemistry)
Anti-SC1 / TCF-19 antibody IHC of human tonsil. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody dilution 5-10 ug/ml.

WB (Western Blot)
Western blot of TCF19/SC1 in 293 cell lysate in the 1) absence and 2) presence of immunizing peptide using at 2 ug/ml. Goat anti-rabbit Ig HRP secondary antibody, and PicoTect ECL substrate solution, were used for this test.



IHC (Immunohistochemistry)
Anti-SC1 / TCF-19 antibody IHC of human brain, cerebellum, white matter. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody dilution 5-10 ug/ml.

ADDITIONAL INFORMATION

Product Categories/Family for anti-TCF19 antibody	Family: Transcription factor
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	117414156
NCBI GeneID	6941
NCBI Accession #	NP_009040.2
NCBI GenBank Nucleotide #	NM_007109.2
UniProt Accession #	Q9Y242 ; Q0EFA8 ; Q13176 ; Q15967 ; Q5SQ89 ; Q5STD6 ; Q5STF5 ; Q9BUM2 ; Q9UBH7 ; A6NCT8 ; B0UY11
Molecular Weight	37,184 Da
NCBI Official Full Name	transcription factor 19
NCBI Official Synonym Full Names	transcription factor 19
NCBI Official Symbol	TCF19
NCBI Official Synonym Symbols	SC1; TCF-19
NCBI Protein Information	transcription factor 19; transcription factor SC1

UniProt Protein Name	Transcription factor 19
UniProt Gene Name	TCF19
UniProt Synonym Gene Names	SC1; TCF-19
UniProt Entry Name	TCF19_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tcf19-tcf-19-antibody-western-blot-immunohistochemistry-elisa/51469>