

Goat anti-Human FASN / Fatty Acid Synthase Polyclonal Antibody | anti-FASN antibody

Goat Polyclonal to Human FASN / Fatty Acid Synthase

CATALOG # AAA51956	HOST Goat	REACTIVITY Human	APPLICATIONS Immunohistochemistry
------------------------------	---------------------	----------------------------	---

PRODUCT SYNONYMS


FASN / Fatty Acid Synthase; Polyclonal Antibody; Goat Polyclonal to Human FASN / Fatty Acid Synthase; Anti-FASN / Fatty Acid Synthase Antibody (Internal) IHC-plus; FASN; Fatty acid synthase; OA-519; SDR27X1; Human FASN; Fatty Acid Synthase; anti-FASN antibody

PRODUCT SPECIFICATIONS


Host	Goat
Reactivity	Human
Clonality	Polyclonal
Specificity	Human FASN / Fatty Acid Synthase.
Purity/Purification	Immunoaffinity Purified
Form/Format	Tris-buffered saline, pH 7.3, 0.5% BSA, 0.02% sodium azide
Concentration	0.5 mg/ml (varies by lot)
Sequence Length	2511
Applicable Applications for anti-FASN antibody	IHC (Immunohistochemistry)
Target	Human FASN / Fatty Acid Synthase
Immunogen	FASN / Fatty Acid Synthase antibody was raised against synthetic peptide YRDRDSQRDLVE from internal region of human FASN (NP_004095.4). Percent identity by BLAST analysis: Human, Gorilla (100%); Coccidioides (83%).
Disclaimer	Due to the highly specific nature of antibodies and antigens, we cannot predict or be held responsible with respect to how this antibody will behave in your systems. Researchers using this antibody should conduct optimization studies to achieve the most optimal result possible for their intended application.

Recommended Immunohistochemistry Protocol	The following protocol is a recommendation only, and AAA Biotech makes no guarantee of the results: Tissue Preparation: Formalin fixation and embedding in paraffin wax. Tissue Sectioning: Make 4-um sections and place on pre-cleaned and charged microscope slides. Heat in a tissue-dryingoven for 45 minutes at 60°C. Deparaffinization: Wash dry slides in 3 changes of xylene - 5 minutes each @ RT Rehydration: Wash slides in 3 changes of 100% alcohol - 3 minutes each @ RT Wash slides in 2 changes of 95% alcohol - 3 minutes each @ RT Wash slides in 1 change of 80% alcohol - 3 minutes @ RT Rinse slides in gentle running distilled water - 5 minutes @ RT Antigen retrieval: Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes Remove from heat and let stand at room temperature in buffer - 20 minutes Rinse in 1X TBS with Tween (TBST) -1 minute @ RT Immunostaining: (Do not allow tissues to dry at any time during the staining procedure) Apply a universal protein block - 20 minutes @ RT Drain protein block from slides, apply diluted primary antibody - 45 minutes @ RT Rinse slides in 1 X TBST - 1 minute @ RT Apply a biotinylated secondary antibody appropriate for the primary antibody - 30 minutes @ RT Rinse slides in 1X TBST -1 minute @ RT Apply alkaline phosphatase streptavidin - 30 minutes @ RT Rinse slides in 1X TBST -1 minute @ RT Apply alkaline phosphatase chromogen substrate - 30 minutes @ RT Wash slides in distilled water - 1 minute @ RT Dehydrate: (This method should only be used if the chromogen substrate is alcohol insoluble (e.g. Vector Red, DAB) Wash slides in 2 changes of 80% alcohol - 1 minute each @ RT Wash slides in 2 changes of 95% alcohol - 1 minute each @ RT Wash slides in 3 changes of 100% alcohol - 1 minute each @ RT Wash slides in 3 changes of xylene - 1 minute each @ RT Apply coverslip
Preparation and Storage	Store at -20°C. Minimize freezing and thawing.

IMAGES



IHC (Immunohistochemistry)
Anti-Fatty Acid Synthase antibody IHC of human brain, cerebellum. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 3.75 ug/ml.



IHC (Immunohistochemistry)
Anti-Fatty Acid Synthase antibody IHC of human brain, cortex. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 3.75 ug/ml.

NCBI AND UNIPROT INFORMATION

NCBI GI #	41872631
NCBI GeneID	2194
NCBI Accession #	NP_004095.4
NCBI GenBank Nucleotide #	NM_004104.4
UniProt Accession #	P49327; Q13479; Q16702; Q4LE83; Q6P4U5; Q6SS02; Q969R1; Q96C68; Q96IT0
Molecular Weight	273,427 Da
NCBI Official Full Name	fatty acid synthase
NCBI Official Synonym Full Names	fatty acid synthase
NCBI Official Symbol	FASN
NCBI Official Synonym Symbols	FAS; OA-519; SDR27X1
NCBI Protein Information	fatty acid synthase; short chain dehydrogenase/reductase family 27X, member 1
UniProt Protein Name	Fatty acid synthase
UniProt Gene Name	FASN
UniProt Synonym Gene Names	FAS
UniProt Entry Name	FAS_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fasn-fatty-acid-synthase-antibody-immunohistochemistry-elisa/51956>