

Rabbit GM2A Polyclonal Antibody | anti-GM2A antibody

Rabbit Polyclonal to Human GM2A

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA52180	Rabbit	Gibbon, Gorilla, Human Predicted Reactivity: Monkey (at least 90% immunogen sequence identity)	Immunohistochemistry


PRODUCT SYNONYMS

GM2A; Polyclonal Antibody; Rabbit Polyclonal to Human GM2A; Anti-GM2A Antibody (C-Terminus) IHC-plus; Ganglioside GM2 activator; GM2 ganglioside activator; GM2-AP; SAP-3; Human GM2A; anti-GM2A antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Gibbon, Gorilla, Human Predicted Reactivity: Monkey (at least 90% immunogen sequence identity)
Clonality	Polyclonal
Specificity	Human GM2A. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Purity/Purification	Immunoaffinity Purified
Form/Format	PBS, 0.1% sodium azide.
Concentration	1 mg/ml (varies by lot)
Sequence Length	193
Applicable Applications for anti-GM2A antibody	IHC (Immunohistochemistry)
Target Species	Human
Immunogen Description	Synthetic 16 amino acid peptide from C-terminus of human GM2A. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon (100%); Monkey (94%); Marmoset, Mouse, Rat, Bovine, Hamster, Elephant, Horse (81%).
Immunogen Type	Synthetic peptide
Immunogen	GM2A antibody was raised against synthetic 16 amino acid peptide from C-terminus of human GM2A. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon (100%); Monkey (94%); Marmoset, Mouse, Rat, Bovine, Hamster, Elephant, Horse (81%).
Antigen Modification	C-Terminus
Preparation and Storage	Short term 4 degree C, long term aliquot and store at -20 degree C, avoid freeze thaw cycles.

IMAGES



IHC (Immunohistochemistry)
Anti-GM2A antibody IHC of human kidney. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



IHC (Immunohistochemistry)
Anti-GM2A antibody IHC of human skin. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.

ADDITIONAL INFORMATION

Related Product Information for anti-GM2A antibody	The protein encoded by this gene is a small glycolipid transport protein which acts as a substrate specific co-factor for the lysosomal enzyme, beta-hexosaminidase A. Beta-hexosaminidase A, together with GM2 ganglioside activator protein, catalyzes the degradation of the ganglioside GM2, and other molecules containing terminal N-acetyl hexosamines. Mutations in this gene result in GM2-gangliosidosis type AB or the AB variant of Tay-Sachs disease.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	39995109
NCBI GeneID	2760
NCBI Accession #	NP_000396.2
NCBI GenBank Nucleotide #	NM_000405.4
UniProt Accession #	P17900; Q14426; Q14428; Q6LBL5; B2R699; D3DQH6
Molecular Weight	20,838 Da
NCBI Official Full Name	ganglioside GM2 activator isoform 1
NCBI Official Synonym Full Names	GM2 ganglioside activator
NCBI Official Symbol	GM2A
NCBI Official Synonym Symbols	SAP-3; GM2-AP
NCBI Protein Information	ganglioside GM2 activator; shingolipid activator protein 3; sphingolipid activator protein 3; cerebroside sulfate activator protein
UniProt Protein Name	Ganglioside GM2 activator
UniProt Gene Name	GM2A
UniProt Synonym Gene Names	SAP-3
UniProt Entry Name	SAP3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.