

Mouse CD38 Monoclonal Antibody | anti-CD38 antibody

CD38

CATALOG # AAA59253	HOST Mouse	REACTIVITY Human, Rabbit
------------------------------	----------------------	------------------------------------

PRODUCT SYNONYMS


CD38; Monoclonal Antibody; anti-CD38 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Rabbit
Clonality	Monoclonal
Isotype	IgG1
Clone Number	SPC32
Form/Format	Anti-CD38 is a mouse monoclonal antibody derived from cell culture supernatant that is concentrated, dialyzed, filter sterilized and diluted in buffer pH 7.5, containing BSA and sodium azide as a preservative.
Control	Tonsil, Lymph Node
Localization	Membranous
Immunogen	Recombinant protein encoding the extracellular domain of human CD38.
Reactivity Note	Paraffin, Frozen
Precautions	1. This product contains <0.1% sodium azide (NaN₃) as a preservative. Ensure proper handling procedures are used with this reagent. 2. Always wear personal protective equipment such as a laboratory coat, goggles, and gloves when handling reagents. 3. Dispose of unused solution with copious amounts of water. 4. Do not ingest reagent. If reagent is ingested, seek medical advice immediately. 5. Avoid contact with eyes. If contact occurs, flush with large quantities of water. 6. Follow safety precautions of the heating device used for epitope retrieval (TintoRetriever Pressure Cooker or similar). 7. For additional safety information refer to Safety Data Sheet for this product. 8. For complete recommendations for handling biological specimens, please refer to the CDC document, "Guidelines for Safe Work Practices in Human and AMDL"
Specimen Preparation	Paraffin sections: The antibody can be used on formalin-fixed paraffin-embedded (FFPE) tissue sections. Ensure tissue undergoes appropriate fixation for best results. Pre-treatment of tissues with heat-induced epitope retrieval (HIER) is recommended using ImmunoDNA Retriever with Citrate, ImmunoDNA Retriever with EDTA, or ImmunoDNA Digester. See reverse side for complete protocol. Tissue should remain hydrated via use of Bio SB Immuno/DNA Washer solutions.
Mounting Protocols	For detailed instructions using biodegradable permanent mounting media such as XyGreen PermaMunter (BSB 0169-0174) or organic solvent based resin such as PermaMunter.
Product Limitations	Due to inherent variability present in immunohistochemical procedures (including fixation time of tissues, dilution factor of antibody, retrieval method utilized, and incubation time), optimal performance should be established through the use of positive and negative controls.
Preparation and Storage	Store at 2-8°C

Abbreviated Immunohistochemical Protocol			
Step	ImmunoDetector AP/HRP	PolyDetector AP/HRP	PolyDetector Plus HRP
Peroxidase/AP Blocker	5 min.	5 min.	5 min.
Primary Antibody	30-60 min.	30-60 min.	30-60 min.
1st Step Detection	10 min.	30-45 min.	15 min.
2nd Step Detection	10 min.	Not Applicable	15 min.
Substrate- Chromogen	5-10 min.	5-10 min.	5-10 min.
Counterstain/Coverslip	Varies	Varies	Varies

IHC (Immunohistochemistry)



IHC (Immunohistochemistry)

Antibody Type	Dilution	Volume/QTY
Prediluted	Ready-To-Use	3.0 ml
Prediluted	Ready-To-Use	7.0 ml
Prediluted	Ready-To-Use	15.0 ml
Concentrated	1:25-1:100	0.1 ml
Concentrated	1:25-1:100	0.5 ml
Concentrated	1:25-1:100	1.0 ml
Control Slides		5

Dilution Info

ADDITIONAL INFORMATION

Related Product Information for anti-CD38 antibody	CD38 is a glycoprotein found on the surface of many immune cells (white blood cells), including CD4+, CD8+, B and natural killer cells. It is a marker of cell activation. The CD38 protein has been connected to HIV infection, Leukemias, Myelomas, solid tumors, Type II Diabetes Mellitus and bone metabolism, as well as some genetically-determined conditions. It has also been used as a prognostic marker in Leukemia. CD38 is highly expressed on thymocytes. It is also expressed by early cells of B and T lineages, NK cells, plasma cells, monocytes and macrophages, and may be detected on cells from Multiple Myeloma, ALL (B and T) and some AML. Monoclonal antibodies to CD38 have been shown to be useful in subtyping of Lymphomas and Leukemias, inhibition of B-lymphopoiesis, detection of plasma cells, protection of B-cells from apoptosis, and as a marker for activated B and T-cell proliferation.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	1911103
NCBI GeneID	952
UniProt Accession #	P28907; O00121; O00122; Q96HY4; Q4FCX6
NCBI Official Full Name	CD38
NCBI Official Synonym Full Names	CD38 molecule
NCBI Official Symbol	CD38
NCBI Official Synonym Symbols	T10
NCBI Protein Information	ADP-ribosyl cyclase 1; cADPr hydrolase 1; CD38 antigen (p45); OTTHUMP00000158633; OTTHUMP00000217743; NAD(+) nucleosidase; cyclic ADP-ribose hydrolase 1; ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase
UniProt Protein Name	ADP-ribosyl cyclase 1
UniProt Gene Name	CD38
UniProt Entry Name	CD38_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd38-antibody-clone-spc32-elisa/59253>