

Mouse anti-Human Surfactant protein D/SP-D Monoclonal Antibody | anti-D/SP-D antibody

Surfactant protein D/SP-D (BSB-162)

CATALOG # AAA59390	HOST Mouse	REACTIVITY Human
------------------------------	----------------------	----------------------------


PRODUCT SYNONYMS

Surfactant protein D/SP-D; Monoclonal Antibody; Surfactant protein D/SP-D (BSB-162); surfactant protein D; anti surfactant; sp-d; spd; anti-sp-d anti-spd; sp d; anti-D/SP-D antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1
Clone Number	BSB-162
Form/Format	Paraffin, Frozen Surfactant Protein is a mouse monoclonal antibody derived from cell culture supernatant that is concentrated, dialyzed, filter sterilized and diluted in buffer pH7.5, containing BSA and sodium azide as a preservative.
Immunogen	Synthetic peptide corresponding to the N-terminus of the human SP-D protein.
Control	Placenta, Lung, Pancreas, Adrenal Gland, Lung Squamous Cell Carcinoma, Lung Adenocarcinoma, Gastric GIST
Localization	Cytoplasmic, Membranous
Preparation and Storage	Store at 2-8 degree C (Control Slides: Store at 20-25 degree C)

IMAGES



IHC (Immunohistochemistry)

Surfactant Protein D Antibody Immunohistochemistry on an FFPE Lung Squamous Cell Carcinoma Tissue

Antibody Type	Dilution	Volume/QT
Prediluted	Ready-To-Use	3.0 ml
Prediluted	Ready-To-Use	7.0 ml
Prediluted	Ready-To-Use	15.0 ml
Concentrated	1:25-1:100	0.1 ml
Concentrated	1:25-1:100	0.5 ml
Concentrated	1:25-1:100	1.0 ml
Control Slides		5

Dilution Info

ADDITIONAL INFORMATION

Related Product Information for anti-D/SP-D antibody	Surfactant protein D, also known as SFTPD or SP-D, is a protein encoded by the SFTPD gene. Pulmonary surfactants are essential to proper respiratory structure and function, consisting of ~90% lipids (mostly phospholipids) and 8-10% surfactant-associated proteins. SP-D is a pattern-recognition molecule in the collectin (collagen-containing C-type lectin) family. In the lungs, it is an opsonin that can agglutinate a range of microbes and enhance their clearance via phagocytosis and super-oxidative burst. It can interfere with allergen-IgE interaction and suppress basophil and mast cell activation. Other surfactant proteins like SP-B and SP-C showed strong immunohistochemical expression in Lung Hyperplasias and Adenomas, suggesting that SP-B and SP-C are related to lung tumorigenesis. SP-D is likely an innate immune surveillance molecule against tumor development. SP-D suppresses Lung Cancer progression via interference with epidermal growth factor signaling, and a truncated recombinant human SP-D may induce apoptosis in Pancreatic Adenocarcinoma via Fas-mediated pathway in a p53-independent manner. Studies Have found low expression of Surfactant protein D antibody expression in Lung, Gastric, and Breast cancers and high expression in different stages and grades of Ovarian cancer. Surfactant protein D antibody expression could be associated with a favorable prognosis in Lung Cancer but unfavorable in non-pulmonary sites such as Gastric, Breast, and Ovarian cancers. SP-D detected in circulating tumor cells of Ovarian Cancer patients suggests SP-D can also be used as a potential biomarker. The purified Spike protein of SARS-CoV-2 bound to Vero but not 293T cells and was itself recognized by SP-D, in the lung alveoli. It suggests that SARS-CoV interacts with innate immune mechanisms in the lung through its S-protein and regulates pulmonary inflammation.
--	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/surfactant-protein-d-sp-d-antibody-clone-bsb-162-elisa/59390>