

Rabbit Beta-Catenin Monoclonal Antibody

Beta-Catenin (RM276)

CATALOG #	HOST	REACTIVITY
AAA59398	Rabbit	HumanPredicted: Mouse, Rat, Sheep, Hamster, Cow, Macaque Monkey, African Green Monkey


PRODUCT SYNONYMS

Beta-Catenin; Monoclonal Antibody; Beta-Catenin (RM276); CTNNB1; sc7963; beta-catenin; b-catenin; b catenin; beta catenin; sc-7963; 7963; betacatenin; anti-beta-catenin; anti beta-catenin; anti-beta catenin; anti-betacatenin; anti beta catenin; anti betacatenin; anti beta-Catenin; anti CTNNB1; anti-beta-Catenin; anti-CTNNB1; catenin cadherin associated protein beta 1 88kDa; catenin beta 1; CTNNB; DKFZp686D02253; FLJ25606; FLJ37923; bcatenin; catenins; PRO2286; anti-Beta-Catenin antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human Predicted: Mouse, Rat, Sheep, Hamster, Cow, Macaque Monkey, African Green Monkey
Clonality	Monoclonal
Isotype	IgG
Clone Number	RM276
Form/Format	Paraffin, Frozen Beta-Catenin is a rabbit monoclonal antibody derived from cell culture supernatant that is concentrated, dialyzed, filter sterilized and diluted in buffer pH7.5, containing BSA and sodium azide as a preservative.
Immunogen	A peptide corresponding to human Beta-Catenin.
Control	Fibromatosis of the Breast and Abdomen, Breast, Abdomen, Colon, Testis, Pancreas
Localization	Nuclear, Cytoplasmic, Membranous
Preparation and Storage	Store at 2-8 degree C (Control Slides: Store at 20-25 degree C)

IMAGES



IHC (Immunohistochemistry)

Beta-Catenin Antibody Immunohistochemistry on an FFPE Salivary Gland Tissue

Antibody Type	Dilution	Volume/QT
Prediluted	Ready-To-Use	3.0 ml
Prediluted	Ready-To-Use	7.0 ml
Prediluted	Ready-To-Use	15.0 ml
Concentrated	1:50-1:200	0.1 ml
Concentrated	1:50-1:200	0.5 ml
Concentrated	1:50-1:200	1.0 ml
Control Slides		5

Dilution Info

ADDITIONAL INFORMATION

Related Product Information for anti-Beta-Catenin antibody	Beta-Catenin is a subunit of the Cadherin protein complex. Cadherins are a type of protein normally expressed on the surface of certain cells. Specifically, Beta-Cateinin is a 92 kDa protein normally found in the cytoplasm of the cell in the sub-membranous location. The Beta-Catenin antibody is associated with E-Cadherin and may be essential for the function of E-Cadherin. Mutations in the Beta-Catenin gene result in the nuclear accumulation of this protein. Nuclear accumulation of this protein has been demonstrated in Fibromatosis lesions of the breast and abdomen, and therefore is useful in differentiating this lesion from other spindle-cell lesions that may occur in these locations.
--	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/beta-catenin-antibody-clone-rm276-elisa/59398>