

Rabbit anti-Human BAP1 Monoclonal Antibody | anti-BAP1 antibody

BAP1 (RBT-BAP1), RMab

CATALOG # AAA59491	HOST Rabbit	REACTIVITY Human
------------------------------	-----------------------	----------------------------


PRODUCT SYNONYMS

BAP1; Monoclonal Antibody; BAP1 (RBT-BAP1); RMab; (RBT-BAP1); TFF3; trefoil factor 3 intestinal; ITF; P1B; TFI; anti-tff3; anti-itf; anti-BAP1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Clone Number	RBT-BAP1
Form/Format	This antibody is derived from cell culture supernatant that is concentrated, dialyzed, filter sterilized and diluted in buffer pH 7.5, containing BSA and sodium azide.
Localization	Nuclear
Intended Use	BAP1 (RBT-BAP1), rabbit monoclonal antibody, is a primary antibody intended for laboratory use by trained laboratory personnel in an immunohistochemical (IHC) assay to qualitatively identify the BAP1 protein by light microscopy in normal and/or pathological formalin-fixed, paraffin-embedded (FFPE) human tissue. The clinical interpretation of any staining or its absence should be performed by a qualified pathologist and complemented by morphological studies using proper controls and evaluated within the context of the patient's clinical history and other tests.

IMAGES



IHC (Immunohistochemistry)

IHC of BAP1 on an FFPE Normal Breast tissue

Antibody Type	Dilution	Volume/QTY
Prediluted	Ready-To-Use	3.0 ml
Prediluted	Ready-To-Use	7.0 ml
Prediluted	Ready-To-Use	15.0 ml
Concentrated	1:25-1:100	0.1 ml
Concentrated	1:25-1:100	0.5 ml
Concentrated	1:25-1:100	1.0 ml
Control Slides		5

Dilution Info

ADDITIONAL INFORMATION

Related Product Information for anti-BAP1 antibody	BAP1 or BRCA1 associated protein-1 (ubiquitin carboxy-terminal hydrolase) is a deubiquitinating enzyme that in humans is encoded by the BAP1 gene. In cancer, BAP1 can function both as a tumor suppressor and as a metastasis suppressor. Exome sequencing identified inactivating mutations in BAP1 in 47% of Uveal melanomas, and BAP1 mutation have been found to be strongly associated with metastasis. BAP1 mutations have been identified in aggressive Mesotheliomas with similar mutations as seen in Melanomas. Mutations in the tumor suppressor gene BAP1 occur in approximately 15% of clear cell renal cell carcinoma cases. Sequencing efforts demonstrated worse outcomes in patients with BAP1 mutated clear cell renal cell carcinoma.
--	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/bap1-antibody-clone-rbt-bap1-elisa/59491>