

Rabbit Histone H4ac Polyclonal Antibody | anti-H4ac antibody

Histone H4ac (pan-acetyl) antibody (pAb)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA59803	Rabbit	Human, Mouse, Wide Range Predicted	Western Blot, Immunofluorescence, Immunocytochemistry, Dot Blot, Chromatin Immunoprecipitation, Immunoprecipitation

PRODUCT SYNONYMS

Histone H4ac; Polyclonal Antibody; Histone H4ac (pan-acetyl) antibody (pAb); anti-H4ac antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Wide Range Predicted
Clonality	Polyclonal
Form/Format	Rabbit serum containing 30% glycerol and 0.035% sodium azide. Sodium azide is highly toxic. For your convenience, an IgG version of this antibody that was purified by Protein A Chromatography is also available.
Applicable Applications for anti-H4ac antibody	WB (Western Blot), IF (Immunofluorescence), ICC (Immunocytochemistry), DB (Dot Blot), ChIP (Chromatin immunoprecipitation)
Conjugation	Acetylated
Immunogen	This Histone H4 pan-acetyl antibody was raised against a peptide containing the amino terminal region of Tetrahymena histone H2A acetylated at multiple lysines. (The immunogen has a high degree of homology to mammalian histone H4, so the antibody recognizes acetylated histone H4 in HeLa extracts).
Abs Residue	Lys
Preparation and Storage	Some products may be shipped at room temperature. This will not affect their stability or performance. Avoid repeated freeze/thaw cycles by aliquoting items into single-use fractions for storage at -20 degree C for up to 2 years. Keep all reagents on ice when not in storage.1ug/ul




DB (Dot Blot)

Histone H4 pan-acetyl antibody (pAb) tested by dot blot analysis. Dot blot analysis was used to confirm the specificity of Histone H4 pan-acetyl pAb for acetyl histone H4. Decreasing amounts of acetylated peptides corresponding to the immunogen, a related sequence, as well as unacetylated histone H4 peptides were spotted onto PVDF and probed with the antibody at a 1:10,000 dilution. Lane 1: AGG[ack]GG[ack]GMG[ack]VGA[ack]RHSC Lane 2: AGG[ack]GG[ack]GG[ack]GG[ack]GG[ack]GGC Lane 3: SGRGKGGKGLC Lane 4: GKGGAKRHRKC




WB (Western Blot)

Histone H4 pan-acetyl antibody (pAb) tested by Western blot. HeLa acid extract (20 ug /lane) probed with Histone H4 pan-acetyl polyclonal antibody (1:20,000 dilution). Lane 1: Untreated cells. Lane 2: Cells treated with sodium butyrate.



IF (Immunofluorescence)

Histone H4 pan-acetyl antibody (pAb) tested by immunofluorescence. Staining of HeLa cells with Histone H4 pan-acetyl pAb (1:1,000 dilution, top panel) and DAPI (middle panel), and a merge of both images (bottom panel).



ChIP (Chromatin Immunoprecipitation)

Histone H4 pan-acetyl antibody (pAb) tested by ChIP analysis. Chromatin IP performed using the ChIP-IT Express Kit and HeLa Chromatin (1.5 x 10⁶ cell equivalents per ChIP) using 2 ul of Histone H4 pan-acetyl pAb or the equivalent amount of rabbit IgG as a negative control. Real time, quantitative PCR (RT-qPCR) was performed on DNA purified from each of the ChIP reactions using a primer pair specific for the indicated gene. Data are presented as Fold Enrichment of the ChIP antibody signal versus the negative control IgG using the ddCT method.

ADDITIONAL INFORMATION

Related Product Information for anti-H4ac antibody

Short Description: Our Histone H4ac (pan-acetyl) antibody (pAb) was raised in a Rabbit host. It has been validated for use in Chromatin Immunoprecipitation, Dot blot, Immunocytochemistry, Immunofluorescence and Western blot; it has been shown to react with Human and Mouse samples, but it is predicted that it will react with a wide range of sample types.

Background: Histone H4 is one of the core components of the nucleosome. The nucleosome is the smallest subunit of chromatin and consists of 147 base pairs of DNA wrapped around an octamer of core histone proteins (two each of Histone H2A, Histone H2B, Histone H3 and Histone H4). Histone H1 is a linker histone, present at the interface between the nucleosome core and DNA entry/exit points; it is responsible for establishing higher-order chromatin structure. Chromatin is subject to a variety of chemical modifications, including post-translational modifications of the histone proteins and the methylation of cytosine residues in the DNA. Reported histone modifications include acetylation, methylation, phosphorylation, ubiquitylation, glycosylation, ADP-ribosylation, carbonylation and SUMOylation; they play a major role in regulating gene expression. Lysine N-epsilon-acetylation is a dynamic, reversible and tightly regulated protein and histone modification that plays a major role in chromatin remodeling and in the regulation of gene expression in various cellular functions. Acetylation of histone H4 occurs at several different lysine positions in the histone tail, and is performed by Histone Acetyltransferases (HATs) such as Hat1 or Gcn5. Acetylation of histones is often associated with transcriptional activation.

NCBI AND UNIPROT INFORMATION

NCBI GI #	28173560
NCBI GeneID	121504
NCBI Accession #	NP_778224; AB_2793201
NCBI GenBank Nucleotide #	NM_175054.2
UniProt Accession #	P62805; P02304; P02305; Q6DRA9; Q6FGB8; Q6NWP7; A2VCL0; Q0VAS5; Q6B823
Molecular Weight	11,367 Da
NCBI Official Full Name	histone H4
NCBI Official Synonym Full Names	histone cluster 4, H4
NCBI Official Symbol	HIST4H4
NCBI Official Synonym Symbols	H4/p; HIST1H4A; HIST1H4B; HIST1H4C; HIST1H4D; HIST1H4E; HIST1H4F; HIST1H4H; HIST1H4I; HIST1H4J; HIST1H4K; HIST1H4L; HIST2H4A; HIST2H4B; MGC24116
NCBI Protein Information	histone H4; histone 4, H4; OTTHUMP00000239323
UniProt Protein Name	Histone H4
UniProt Gene Name	HIST1H4A
UniProt Synonym Gene Names	H4/A; H4FA
UniProt Entry Name	H4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/histone-h4ac-antibody-western-blot-immunofluorescence-immunocytochemistry-dot-blot-chromatin-immunoprecipitation-elisa/59803>