

Rabbit Histone H4K16ac Polyclonal Antibody | anti-H4K16ac antibody

Histone H4K16ac antibody (pAb)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA59952	Rabbit	Budding Yeast, Drosophila, Human, Wide Range Predicted	Western Blot, Immunohistochemistry, Dot Blot, Chromatin Immunoprecipitation, Immunoprecipitation

PRODUCT SYNONYMS

Histone H4K16ac; Polyclonal Antibody; Histone H4K16ac antibody (pAb); anti-H4K16ac antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Budding Yeast, Drosophila, Human, Wide Range Predicted
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Protein A Chromatography
Form/Format	Purified IgG in PBS (pH7.5) with 30% glycerol and 0.035% sodium azide. Sodium azide is highly toxic. For your convenience, an unpurified serum version of this antibody is also available.
Concentration	1ug/ul (varies by lot)
Applicable Applications for anti-H4K16ac antibody	WB (Western Blot), IHC (Immunohistochemistry), DB (Dot Blot), ChIP (Chromatin immunoprecipitation)
Conjugation	Acetylated
Immunogen	This Histone H4 acetyl Lys16 antibody was raised against a peptide including acetyl-lysine 16 of histone H4.
Abs Residue	Lys
Preparation and Storage	Some products may be shipped at room temperature. This will not affect their stability or performance. Avoid repeated freeze/thaw cycles by aliquoting items into single-use fractions for storage at -20 degree C for up to 2 years. Keep all reagents on ice when not in storage.1ug/ul

IMAGES




DB (Dot Blot)

Histone H4 acetyl Lys16 antibody (pAb) tested by dot blot analysis. Dot blot analysis was used to confirm the specificity of Histone H4 acetyl Lys16 antibody for acetyl Lys16 histone H4. Acetylated peptides corresponding to the immunogen and related peptides were spotted onto PVDF and probed with Histone H4 acetyl Lys16 antibody at a dilution of 1 ug/ml. The amount of peptide (picomoles) spotted is indicated next to each row. Lane 1: acetyl-Lys5 peptide. Lane 2: unmodified Lys5 peptide. Lane 3: acetyl-Lys8 peptide. Lane 4: unmodified Lys8 peptide. Lane 5: acetyl-Lys12 peptide. Lane 6: unmodified Lys12 peptide. Lane 7: acetyl-Lys16 peptide. Lane 8: unmodified Lys16 peptide.



WB (Western Blot)

Histone H4 acetyl Lys16 antibody (pAb) tested by Western blot. HeLa nuclear extract (20 ug per lane) probed with Histone H4 acetyl Lys16 antibody (1 ug per ml). Lane 1: no treatment. Lane 2: cells treated with sodium butyrate.



ChIP (Chromatin Immunoprecipitation)

Histone H4 acetyl Lys16 antibody tested by ChIP analysis. Chromatin IP performed using the ChIP-IT Express Kit and HeLa Chromatin (1.5 x 10⁶ cell equivalents per ChIP) using 10 ug of Histone H4 acetyl Lys16 pAb or the equivalent amount of rabbit IgG as a negative control. Real time, quantitative PCR (RT-qPCR) was performed on DNA purified from each of the ChIP reactions using a primer pair specific for the indicated gene. Data are presented as Fold Enrichment of the ChIP antibody signal versus the negative control IgG using the ddCT method.

ADDITIONAL INFORMATION

Related Product Information for anti-H4K16ac antibody	Short Description: Our Histone H4K16ac antibody (pAb) was raised in a Rabbit host. It has been validated for use in Chromatin Immunoprecipitation, Dot blot, Immunohistochemistry and Western blot; it has been shown to react with Budding Yeast, Drosophila and Human samples, but it is predicted that it will react with a wide range of sample types. Background: Histone H4 is one of the core components of the nucleosome. The nucleosome is the smallest subunit of chromatin and consists of 147 base pairs of DNA wrapped around an octamer of core histone proteins (two each of Histone H2A, Histone H2B, Histone H3 and Histone H4). Histone H1 is a linker histone, present at the interface between the nucleosome core and DNA entry/exit points; it is responsible for establishing higher-order chromatin structure. Chromatin is subject to a variety of chemical modifications, including post-translational modifications of the histone proteins and the methylation of cytosine residues in the DNA. Reported histone modifications include acetylation, methylation, phosphorylation, ubiquitylation, glycosylation, ADP-ribosylation, carbonylation and SUMOylation; they play a major role in regulating gene expression. Lysine N-epsilon-acetylation is a dynamic, reversible and tightly regulated protein and histone modification that plays a major role in chromatin remodeling and in the regulation of gene expression in various cellular functions. NoRC is a SMARCA5 (SNF2h)-containing chromatin remodeling complex. The bromodomain of TIP5, the large subunit of NoRC, interacts with acetylated Histone H4 Lys16, (H4K16ac) and cooperates with an adjacent PHD finger to recruit histone deacetylases and DNA methyltransferases to rDNA, leading to the silencing of rDNA.
Product Categories/Family for anti-H4K16ac antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	28173560
NCBI GeneID	121504
NCBI Accession #	NP_778224; AB_2753164
NCBI GenBank Nucleotide #	NM_175054.2
UniProt Accession #	P62805; P02304; P02305; Q6DRA9; Q6FGB8; Q6NWP7; A2VCL0; Q0VAS5; Q6B823
Molecular Weight	11,367 Da
NCBI Official Full Name	histone H4
NCBI Official Synonym Full Names	histone cluster 4, H4
NCBI Official Symbol	HIST4H4
NCBI Official Synonym Symbols	H4/p; HIST1H4A; HIST1H4B; HIST1H4C; HIST1H4D; HIST1H4E; HIST1H4F; HIST1H4H; HIST1H4I; HIST1H4J; HIST1H4K; HIST1H4L; HIST2H4A; HIST2H4B; MGC24116
NCBI Protein Information	histone H4; histone 4, H4; OTTHUMP00000239323
UniProt Protein Name	Histone H4
UniProt Gene Name	HIST1H4A
UniProt Synonym Gene Names	H4/A; H4FA
UniProt Entry Name	H4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/histone-h4k16ac-antibody-western-blot-immunohistochemistry-dot-blot-chromatin-immunoprecipitation-elisa/59952>