

Rabbit anti-Human MED1 Polyclonal Antibody | anti-MED1 antibody

MED1 antibody (pAb)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA59972	Rabbit	Human	Western Blot, Immunoprecipitation


PRODUCT SYNONYMS

MED1; Polyclonal Antibody; MED1 antibody (pAb); PBP; CRSP1; RB18A; TRIP2; PPARBP; CRSP200; DRIP205; DRIP230; PPARGBP; TRAP220; anti-MED1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purified
Form/Format	Purified IgG in 70mM Tris (pH8), 105mM NaCl, 31mM glycine, 0.07mM EDTA, 30% glycerol and 0.035% sodium azide. Sodium azide is highly toxic.
Applicable Applications for anti-MED1 antibody	WB (Western Blot), IP (Immunoprecipitation)
Immunogen	This MED1 antibody was raised against a peptide within the C-terminal region of human MED1.
Preparation and Storage	Some products may be shipped at room temperature. This will not affect their stability or performance. Avoid repeated freeze/thaw cycles by aliquoting items into single-use fractions for storage at -20 degree C for up to 2 years. Keep all reagents on ice when not in storage.1ug/ul

IMAGES



WB (Western Blot)

MED1 antibody (pAb) tested by Western blot. Nuclear extract (30 ug) of HeLa cells probed with MED1 antibody at a dilution of 1:2,000.



WB (Western Blot)

MED1 antibody (pAb) tested by Immunoprecipitation. 10 ul of MED1 antibody was used to immunoprecipitate MED1 from 250 ug of HeLa nuclear cell extract (lane 2). 10 ul of rabbit IgG was used as a negative control (lane 1). The immunoprecipitated protein was detected by Western blotting using the MED1 antibody at a dilution of 1:2,000.

ADDITIONAL INFORMATION

Related Product Information for anti-MED1 antibody	Short Description: Our MED1 antibody (pAb) was raised in a Rabbit host. It has been validated for use in Immunoprecipitation and Western blot; it has been shown to react with Human samples. Background: MED1 (Mediator complex subunit 1, TRIP2, Thyroid receptor-interacting protein 2) is a component of the Mediator complex, a coactivator involved in the regulated transcription of most genes transcribed by RNA polymerase II. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional pre-initiation complex with RNA polymerase II and the general transcription factors. MED1 is involved in supporting hematopoietic stem and/or progenitor cell function.
Product Categories/Family for anti-MED1 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

Molecular Weight	210kDa
------------------	--------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/med1-antibody-western-blot-immunoprecipitation-elisa/59972>