

Rabbit anti-Human MED15 Polyclonal Antibody | anti-MED15 antibody

MED15 antibody (pAb)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA60052	Rabbit	Human	Western Blot, Immunoprecipitation


PRODUCT SYNONYMS

MED15; Polyclonal Antibody; MED15 antibody (pAb); TIG1; CAG7A; CTG7A; PCQAP; TIG-1; TNRC7; ARC105; anti-MED15 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purified
Form/Format	Purified IgG in PBS with 30% glycerol and 0.035% sodium azide. Sodium azide is highly toxic.
Applicable Applications for anti-MED15 antibody	WB (Western Blot), IP (Immunoprecipitation)
Immunogen	This antibody was raised against a peptide within the C-terminal region of human MED15.
Preparation and Storage	Some products may be shipped at room temperature. This will not affect their stability or performance. Avoid repeated freeze/thaw cycles by aliquoting items into single-use fractions for storage at -20 degree C for up to 2 years. Keep all reagents on ice when not in storage.1ug/ul

IMAGES



WB (Western Blot)

MED15 antibody (pAb) tested by Western blot. HeLa whole-cell extract (20 ug) probed with MED15 antibody at a dilution of 1:1,000.



WB (Western Blot)

MED15 antibody (pAb) tested by Immunoprecipitation. 10 ul of MED15 antibody was used to immunoprecipitate MED15 from 400 ug of HeLa whole cell extract (lane 2). 10 ul of rabbit IgG was used as a negative control (lane 1). The immunoprecipitated protein was detected by Western blotting using the MED15 antibody at a dilution of 1:1,000.

ADDITIONAL INFORMATION

Related Product Information for anti-MED15 antibody	Short Description: Our MED15 antibody (pAb) was raised in a Rabbit host. It has been validated for use in Immunoprecipitation and Western blot; it has been shown to react with Human samples. Background: MED15 (mediator of RNA polymerase II transcription subunit 15) is a component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. Required for cholesterol-dependent gene regulation. Positively regulates the Nodal signaling pathway.
Product Categories/Family for anti-MED15 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

Molecular Weight	100kDa
------------------	--------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/med15-antibody-western-blot-immunoprecipitation-elisa/60052>