

IGF2BP3 recombinant protein

Recombinant IGF2BP3 protein

CATALOG #
AAA60211

HOST
Baculovirus


PRODUCT SYNONYMS

IGF2BP3; Recombinant IGF2BP3 protein; VICKZ3; KOC1; IMP3; Insulin-like growth factor 2 mRNA-binding protein 3; IGF2BP3 recombinant protein

PRODUCT SPECIFICATIONS


Host	Baculovirus
Form/Format	Recombinant IGF2BP3 protein is supplied in 25mM HEPES-NaOH pH7.5, 300mM NaCl, 10% glycerol, 0.04% Triton X-100, and 0.5mM TCEP.
Protein Species	Human
Tag	DYKDDDDK-Tag
Notes	This product was manufactured as described in Protein Details. Where possible, We has developed functional or activity assays for recombinant proteins. Additional characterization such as enzyme kinetic activity assays, inhibitor screening or other biological activity assays may not have been performed for every product. All available data for a given product is shown on the lot-specific Technical Data Sheet.
Dry Ice Note	The product may be shipped with dry ice, and an additional fee may be added to your shipping cost.
Preparation and Storage	Recombinant proteins in solution are temperature sensitive and must be stored at -80 degree C to prevent degradation. Avoid repeated freeze/thaw cycles and keep on ice when not in storage. Shipping Temp: Dry Ice

IMAGES




Application Data

HTRF for IGF2BP3 activity 1 uM ACTB zipcode ssDNA (1212-1313) was incubated with different concentrations of IGF2BP3 protein in a 10 ul reaction system containing 50 mM HEPES-NaOH pH 7.5, 0.1% BSA for 1 hour, then 10 ul anti-DYKDDDDK antibody and SA-XL665 mixture (1:100 dilution in the same buffer) was added to each reaction system and incubated for 30 min. All the operations and reactions were performed at room temperature. HTRF assay was used for detection.



Application Data

HTRF for IGF2BP3 activity 1 uM ACTB zipcode ssDNA (1212-1313) was incubated with different concentrations of IGF2BP3 protein in a 10 ul reaction system containing 50 mM HEPES-NaOH pH 7.5, 0.1% BSA for 1 hour, then 10 ul anti-DYKDDDDK antibody and SA-XL665 mixture (1:100 dilution in the same buffer) was added to each reaction system and incubated for 30 min. All the operations and reactions were performed at room temperature. HTRF assay was used for detection.



SDS-PAGE

Recombinant IGF2BP3 protein gel 10% SDS-PAGE with Coomassie blue staining MW: 65 kDa Purity: >85%

ADDITIONAL INFORMATION

Related Product Information for IGF2BP3 recombinant protein	Short Description: Full-length human IGF2BP3 protein (accession number NP_006538.2) was expressed in a baculovirus system with an N-terminal DYKDDDDK-Tag. The molecular weight of the protein is 65 kDa. Background: IGF2BP3 (Insulin Like Growth Factor 2 mRNA Binding Protein 3), also called as VICKZ3, IMP3 or KOC1, is a member of the insulin-like growth factor 2 mRNA-binding protein family. The mammalian IGF2 mRNA binding proteins (IGF2BPs) family encompasses three RNA-binding proteins (RBPs) controlling the cytoplasmic fate of mRNAs in development, somatic cells and human diseases. IGF2BPs are RNA-binding factors that recruit target transcripts to cytoplasmic protein-RNA complexes (mRNPs). This transcript 'caging' into mRNPs allows mRNA transport and transient storage. It also modulates the rate and location at which target transcripts encounter the translational apparatus and shields them from endonuclease attacks or microRNA-mediated degradation. It functions by binding to the mRNAs of certain genes, including insulin-like growth factor 2, beta-actin and beta-transducin repeat-containing protein, and regulating their translation. IGF2BP3 has been reported to be involved in the progression of various cancers, including those in the pancreas, colon or rectum, lungs, and ovaries. According to a study, it is associated with aggressive tumor features and unfavorable outcomes. Recent studies revealed that the association of IGF2BPs with target mRNAs, e.g. the MYC mRNA, is enhanced by the N6-methyladenosine (m6A) modification of target transcripts suggesting IGF2BPs as novel m6Areaders. N6-methyladenosine (m6A) is the most prevalent modification in eukaryotic RNAs. The biological importance of m6A relies on m6A readers, which control mRNA fate and function.
Product Categories/Family for IGF2BP3 recombinant protein	Recombinant Proteins ; Growth Factor

NCBI AND UNIPROT INFORMATION

NCBI GI #	30795212
NCBI GeneID	10643
NCBI Accession #	NP_006538.2
NCBI GenBank Nucleotide #	NM_006547.2
UniProt Accession #	O00425 ; Q63HM0 ; Q6MZZ2 ; Q86VB1 ; A0A4Z5
Molecular Weight	63,705 Da
NCBI Official Full Name	insulin-like growth factor 2 mRNA-binding protein 3
NCBI Official Synonym Full Names	insulin-like growth factor 2 mRNA binding protein 3
NCBI Official Symbol	IGF2BP3
NCBI Official Synonym Symbols	CT98; IMP3; KOC1; IMP-3; VICKZ3

NCBI Protein Information	insulin-like growth factor 2 mRNA-binding protein 3; hKOC; VICKZ family member 3; cancer/testis antigen 98; IGF2 mRNA-binding protein 3; IGF II mRNA binding protein 3; IGF-II mRNA-binding protein 3; KH domain containing protein overexpressed in cancer; KH
UniProt Protein Name	Insulin-like growth factor 2 mRNA-binding protein 3
UniProt Gene Name	IGF2BP3
UniProt Synonym Gene Names	IMP3; KOC1; VICKZ3; IGF2 mRNA-binding protein 3; IMP-3; hKOC
UniProt Entry Name	IF2B3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/igf2bp3-recombinant-protein-baculovirus/60211>