

Goat BCOR/BCL6 co-repressor Polyclonal Antibody | anti-BCOR antibody

Goat anti-BCOR/BCL6 co-repressor Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA61006	Goat	Tested: Human; Expected from sequence similarity: Human, Mouse, Dog, Pig, Cow	Immunofluorescence, ELISA


PRODUCT SYNONYMS

BCOR/BCL6 co-repressor; Polyclonal Antibody; Goat anti-BCOR/BCL6 co-repressor Antibody; BCOR; BCL-6 interacting corepressor; FLJ20285; KIAA1575; BCL6 co-repressor; ANOP2; FLJ38041; MAA2; MGC131961; MGC71031; 5830466J11Rik; 8430401K06Rik; MCOPS2; BCOR antibody; BCL-6 interacting corepressor antibody; FLJ20285 antibody; KIAA1575 antibody; BCL6 co-repressor antibody; ANOP2 antibody; FLJ38041 antibody; MAA2 antibody; MGC131961 antibody; MGC71031 antibody; 5830466J11Rik antibody; 8430401K06Rik antibody; MCOPS2 antibody; BCOR / BCL6 co-repressor; anti-BCOR antibody


PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Tested: Human; Expected from sequence similarity: Human, Mouse, Dog, Pig, Cow
Clonality	Polyclonal
Purity/Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Form/Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Sequence	LSATPLYGNV HSW
Sequence Length	1721
Applicable Applications for anti-BCOR antibody	IF (Immunofluorescence), ELISA (Peptide ELISA)
Immunogen	Peptide with sequence LSATPLYGNVHSW-C, from the N Terminus of the protein sequence according to NP_060215.4; NP_001116855.1; NP_001116856.1; NP_001116857.1.
Epitope	N Terminus
Note	This antibody is expected to recognise all reported isoforms. Variants (NP_060215.4; NP_001116855.1) encode the same isoform.
Preparation and Storage	Aliquot and store at -20 degree C. Minimize freezing and thawing.

IMAGES



IF (Immunofluorescence)
Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and weak cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



IF (Immunofluorescence)
Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

NCBI AND UNIPROT INFORMATION

NCBI GI #	21071037
NCBI GeneID	54880
NCBI Accession #	NP_060215.4; NP_001116855.1; NP_001116856.1; NP_001116857.1
NCBI GenBank Nucleotide #	NM_017745.5
NCBI Official Full Name	BCL-6 corepressor isoform a
NCBI Official Synonym Full Names	BCL6 corepressor
NCBI Official Symbol	BCOR
NCBI Official Synonym Symbols	MAA2; ANOP2; MCOPS2
NCBI Protein Information	BCL-6 corepressor
UniProt Protein Name	BCL-6 corepressor
UniProt Gene Name	BCOR
UniProt Synonym Gene Names	KIAA1575; BCoR
UniProt Entry Name	BCOR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/bcor-bcl6-co-repressor-antibody-immunofluorescence-peptide-elisa-elisa/61006>