

Goat B5 Receptor/PCID1/EIF3M Polyclonal Antibody | anti-EIF3M antibody

Goat anti-B5 Receptor/PCID1/EIF3M Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA61177	Goat	Tested: Human; Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

B5 Receptor/PCID1/EIF3M; Polyclonal Antibody; Goat anti-B5 Receptor/PCID1/EIF3M Antibody; EIF3M; B5 receptor; PCID1; PCI domain containing 1 (herpesvirus entry mediator); FLJ29030; GA17; hfl-B5; dendritic cell protein; fetal lung protein B5; eukaryotic translation initiation factor 3; subunit M; B5; B5 receptor antibody; PCID1 antibody; PCI domain containing 1 (herpesvirus entry mediator) antibody; FLJ29030 antibody; GA17 antibody; hfl-B5 antibody; dendritic cell protein antibody; fetal lung protein B5 antibody; EIF3M antibody; subunit M antibody; B5 antibody; B5 Receptor; B5 Receptor / PCID1 / EIF3M; anti-EIF3M antibody

PRODUCT SPECIFICATIONS


Host	Goat
Reactivity	Tested: Human; Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow
Clonality	Polyclonal
Purity/Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Form/Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Concentration	100ug specific antibody in 200ul (varies by lot)
Sequence	NAWKQNLNKV KN
Sequence Length	374
Applicable Applications for anti-EIF3M antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA (Peptide ELISA)
Immunogen	Peptide with sequence C-NAWKQNLNKVKN, from the C Terminus of the protein sequence according to NP_006351.2.
Epitope	C Terminus
Preparation and Storage	Aliquot and store at -20 degree C. Minimize freezing and thawing.

IMAGES



IHC (Immunohistochemistry)

5ug/ml staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



WB (Western Blot)

0.1ug/ml staining of A549 lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

NCBI AND UNIPROT INFORMATION

NCBI GI #	23397429
NCBI GeneID	10480

NCBI Accession #	NP_006351.2
NCBI GenBank Nucleotide #	NM_006360.5
Molecular Weight	27,865 Da
NCBI Official Full Name	eukaryotic translation initiation factor 3 subunit M isoform 1
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 3 subunit M
NCBI Official Symbol	EIF3M
NCBI Official Synonym Symbols	B5; GA17; PCID1; TANGO7; hfl-B5
NCBI Protein Information	eukaryotic translation initiation factor 3 subunit M
UniProt Protein Name	Eukaryotic translation initiation factor 3 subunit M
UniProt Gene Name	EIF3M
UniProt Synonym Gene Names	eIF3m; hFL-B5
UniProt Entry Name	EIF3M_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/b5-receptor-pcid1-eif3m-antibody-immunohistochemistry-western-blot-peptide-elisa-elisa/61177>