

Goat TAp63alpha/deltaNp63alpha Polyclonal Antibody | anti-TP63 antibody

Goat anti-TAp63alpha/deltaNp63alpha Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA61559	Goat	Tested: Human, Mouse, Rat; Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow	Western Blot, ELISA

PRODUCT SYNONYMS

TAp63alpha/deltaNp63alpha; Polyclonal Antibody; Goat anti-TAp63alpha/deltaNp63alpha Antibody; Goat Anti-Trp63 (mouse) Antibody; TP63; tumor protein p63; AIS; B(p51A); B(p51B); EEC3; KET; LMS; NBP; OFC8; RHS; SHFM4; TP53CP; TP53L; TP73L; p40; p51; p53CP; p63; p73H; p73L; CUSP; OTTHUMP00000209734; OTTHUMP00000209738; OTTHUMP00000209740; OTTHUMP00000209741; OTTHUMP00000209742; amplif; TP63 antibody; tumor protein p63 antibody; AIS antibody; B(p51A) antibody; B(p51B) antibody; EEC3 antibody; KET antibody; LMS antibody; NBP antibody; RHS antibody; SHFM4 antibody; TP53CP antibody; TP53L antibody; TP73L antibody; p40 antibody; p51 antibody; p53CP antibody; p63 antibody; p73H antibody; p73L antibody; CUSP antibody; TAp63alpha; deltaNp63alpha; TAp63alpha / deltaNp63alpha; anti-TP63 antibody


PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Tested: Human, Mouse, Rat; Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow
Clonality	Polyclonal
Purity/Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Form/Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Concentration	100ug specific antibody in 200ul (varies by lot)
Sequence	RNKQQRKEE GE
Sequence Length	680
Applicable Applications for anti-TP63 antibody	WB (Western Blot), ELISA (Peptide ELISA)
Immunogen	Peptide with sequence C-RNKQQRKEEGE, from the C Terminus of the protein sequence according to NP_003713.3; NP_001108452.1.
Epitope	C Terminus
Note	This antibody is expected to recognize isoform 1 (NP_003713.3) and isoform 4 (NP_001108452.1) of TP63.
Preparation and Storage	Aliquot and store at -20 degree C. Minimize freezing and thawing.

IMAGES



WB (Western Blot)
 0.3ug/ml staining of Mouse (A) and Rat (B) Heart lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



WB (Western Blot)
 0.3ug/ml staining of Human Heart lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

NCBI AND UNIPROT INFORMATION

NCBI GI #	31543818
-----------	----------

NCBI GeneID	8626
NCBI Accession #	NP_003713.3; NP_001108452.1
NCBI GenBank Nucleotide #	NM_003722.4
Molecular Weight	65,288 Da
NCBI Official Full Name	tumor protein 63 isoform 1
NCBI Official Synonym Full Names	tumor protein p63
NCBI Official Symbol	TP63
NCBI Official Synonym Symbols	AIS; KET; LMS; NBP; RHS; p40; p51; p63; EEC3; OFC8; p73H; p73L; SHFM4; TP53L; TP73L; p53CP; TP53CP; B(p51A); B(p51B)
NCBI Protein Information	tumor protein 63
UniProt Protein Name	Tumor protein 63
UniProt Gene Name	TP63
UniProt Synonym Gene Names	KET; P63; P73H; P73L; TP73L; p63; CUSP; TP63; p73L
UniProt Entry Name	P63_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tap63alpha-deltanp63alpha-antibody-western-blot-peptide-elisa-elisa/61559>