

Goat Terminal Uridylyltransferase 4 Polyclonal Antibody | anti-Zcchc11 antibody

Goat anti-Terminal Uridylyltransferase 4 (mouse) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA61599	Goat	Tested: Mouse, Rat; Expected from sequence similarity: Mouse, Rat	Western Blot, ELISA


PRODUCT SYNONYMS

Terminal Uridylyltransferase 4; Polyclonal Antibody; Goat anti-Terminal Uridylyltransferase 4 (mouse) Antibody; Zcchc11; zinc finger; CCHC domain containing 11; RP23-406B13.1; 6030404K05Rik; 9230115F04Rik; KIAA0191; PPAPD3; mKIAA0191; OTTMUSP00000008601; TUTase 4; terminal uridylyltransferase 4; zinc finger CCHC domain-containing protein 11; 6030404K05Rik antibody; 9230115F04Rik antibody; KIAA0191 antibody; mKIAA0191 antibody; OTTMUSP00000008601 antibody; PPAPD3 antibody; RP23-406B13.1 antibody; terminal uridylyltransferase 4 antibody; TUTase 4 antibody; zinc finger CCHC domain-containing protein 11 antibody; CCHC domain containing 11 antibody; Zcchc11 antibody; Terminal Uridylyltransferase 4 (mouse); anti-Zcchc11 antibody


PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Tested: Mouse, Rat; Expected from sequence similarity: Mouse, Rat
Clonality	Polyclonal
Purity/Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Form/Format	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Concentration	100ug specific antibody in 200ul (varies by lot)
Sequence	QKSAEKQGDLPSET
Sequence Length	1644
Applicable Applications for anti-Zcchc11 antibody	WB (Western Blot), ELISA (Peptide ELISA)
Immunogen	Peptide with sequence C-QKSAEKQGDLPSET, from the internal region of the protein sequence according to NP_780681.2.
Epitope	Internal region
Preparation and Storage	Aliquot and store at -20 degree C. Minimize freezing and thawing.

IMAGES



WB (Western Blot)
0.2ug/ml) staining of different Testis lysates from wildtype 8-9 week old C57BL/6 (wt) and knock-out (KO) mice (35ug protein in RIPA buffer). Primary incubation was 2 hours. Detected by chemiluminescence. Data obtained from an anonymous customer.



WB (Western Blot)
0. 05ug/ml) staining of Rat Brain lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

NCBI AND UNIPROT INFORMATION

NCBI GI #	83977461
NCBI GeneID	230594
NCBI Accession #	NP_780681.2
NCBI GenBank Nucleotide #	NM_175472.3
Molecular Weight	184,650 Da
NCBI Official Full Name	terminal uridylyltransferase 4
NCBI Official Synonym Full Names	zinc finger, CCHC domain containing 11
NCBI Official Symbol	Zcchc11
NCBI Official Synonym Symbols	PPAPD3; mKIAA0191; 6030404K05Rik; 9230115F04Rik
NCBI Protein Information	terminal uridylyltransferase 4
UniProt Protein Name	Terminal uridylyltransferase 4
UniProt Gene Name	Zcchc11
UniProt Synonym Gene Names	Kiaa0191; Tut4; TUTase 4
UniProt Entry Name	TUT4_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/terminal-uridylyltransferase-4-antibody-western-blot-peptide-elisa-elisa/61599>