

Mouse p21WAF1 Monoclonal Antibody | anti-CDKN1A antibody

p21WAF1 (Tumor Suppressor Protein) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62508	Mouse	Human, Chimpanzee, Monkey, Mouse, Rat	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

p21WAF1; Monoclonal Antibody; p21WAF1 (Tumor Suppressor Protein) Mouse Monoclonal Antibody; Activating Fragment 1; CAP20; CDK-interacting protein 1; CDKI; CDKN1; CDKN1A; CIP1; Cyclin-dependent kinase inhibitor 1A (p21; Cip1); DNA Synthesis Inhibitor; MDA6; Melanoma Differentiation Associated Protein 6; p21Cip1/Waf1; PIC1; SDI1; SLC12A9; Wild type p53 activated fragment 1 (WAF1); anti-CDKN1A antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Chimpanzee, Monkey, Mouse, Rat
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	WA-1
Specificity	This MAb recognizes a 21kDa protein, identified as the p21WAF1 tumor suppressor protein. This MAb is highly specific to p21 and shows no cross-reaction with other closely related mitotic inhibitors. p21WAF1 is a specific inhibitor of cdk's and a tumor suppressor involved in the pathogenesis of a variety of malignancies. The expression of this gene acts as an inhibitor of the cell cycle during G1 phase and is tightly controlled by the tumor suppressor protein p53. Its expression is induced by the wild type, but not mutant, p53 suppressor protein. Normal cells generally display a rather intense nuclear p21 expression. Loss of p21 expression has been reported in many carcinomas (gastric carcinoma, non-small cell lung carcinoma, thyroid carcinoma). In ELISA, MAb WA-1 is useful either as a solid phase or for detection of p21 protein.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	164
Applicable Applications for anti-CDKN1A antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Nuclear
Immunogen	Human recombinant p21 protein
Hu-Chromosome Location	6p21.31
Positive Control	HeLa Cells. Skin, colon, or breast carcinoma.
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Bladder Carcinoma stained with p21 Monoclonal Antibody (WA-1).

ADDITIONAL INFORMATION

References	Kovacic, J. et al., Int.J.Oncol. 9(suppl.), 835 (1996).
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	11386203
NCBI GeneID	1026
NCBI Accession #	NP_000380.1
NCBI GenBank Nucleotide #	NM_000389.4
UniProt Accession #	P38936; Q14010; Q6FI05; Q9BUT4
Molecular Weight	21kDa
NCBI Official Full Name	cyclin-dependent kinase inhibitor 1 isoform 1
NCBI Official Synonym Full Names	cyclin dependent kinase inhibitor 1A
NCBI Official Symbol	CDKN1A
NCBI Official Synonym Symbols	P21; CIP1; SDI1; WAF1; CAP20; CDKN1; MDA-6; p21CIP1
NCBI Protein Information	cyclin-dependent kinase inhibitor 1
UniProt Protein Name	Cyclin-dependent kinase inhibitor 1
UniProt Gene Name	CDKN1A
UniProt Synonym Gene Names	CAP20; CDKN1; CIP1; MDA6; PIC1; SDI1; WAF1; MDA-6
UniProt Entry Name	CDN1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.