

Mouse anti-Human Cytokeratin 8/18 Monoclonal Antibody | anti-KRT8 antibody

Cytokeratin 8/18 Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62519	Mouse	Human	Immunohistochemistry, Western Blot, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Cytokeratin 8/18; Monoclonal Antibody; Cytokeratin 8/18 Mouse Monoclonal Antibody; CARD2; CK8; CYK8; CYKER; Cytokeratin Endo A; DreK8; EndoA; K2C8; K8; Keratin 8; Krt 2.8; KRT8; Type-II Keratin Kb8. Cell Proliferation-inducing Gene 46 Protein; CK18; CYK18 Cytokeratin Endo B; K18; Keratin-18; Kerd; KRT18; anti-KRT8 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Clone Number	K8.8 + DC10
Specificity	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This MAb cocktail recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the normal epidermis so only Paget cells are stained. Immunohistochemical staining with this MAb is indistinguishable from that obtained with monoclonal antibody 5D3.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	483
Applicable Applications for anti-KRT8 antibody	IHC (Immunohistochemistry), WB (Western Blot), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Cytoplasmic
Immunogen	Keratin preparation from a human carcinoma (K8.8); PMC-42 human breast carcinoma cells (DC10)
Hu-Chromosome Location	17q21.2 (CK8); 12q13.13 (CK18)
Positive Control	MCF-7 or A431 cells. Skin, Colon, lung or breast carcinoma
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with Cytokeratin 8/18 Monoclonal Antibody (K8.8 + DC10).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Ovarian Cancer stained with Cytokeratin 8/18 Monoclonal Antibody (K8.8 + DC10).

ADDITIONAL INFORMATION

References	Angus B et. al. J Path, 153:377-384, 1987. Angus B et. al. J Path, 155:71-75, 198
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	372466577
NCBI GeneID	3856
NCBI Accession #	NP_001243222.1
NCBI GenBank Nucleotide #	NM_001256293.1
UniProt Accession #	P05787; Q14099; Q14716; Q14717; Q53GJ0; Q6DHW5; Q6GMY0; Q6P4C7; Q96J60; A8K4H3; B0AZN5; F8VXB4
Molecular Weight	52.5kDa (CK8); 45kDa (CK18)
NCBI Official Full Name	keratin, type II cytoskeletal 8 isoform 2
NCBI Official Synonym Full Names	keratin 8
NCBI Official Symbol	KRT8
NCBI Official Synonym Symbols	K8; KO; CK8; CK-8; CYK8; K2C8; CARD2
NCBI Protein Information	keratin, type II cytoskeletal 8
UniProt Protein Name	Keratin, type II cytoskeletal 8
UniProt Gene Name	KRT8
UniProt Synonym Gene Names	CYK8; CK-8; K8
UniProt Entry Name	K2C8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cytokeratin-8-18-antibody-clone-k8-8-dc10-immunohistochemistry-western-blot-immunofluorescence-flow-cytometry-elisa/62519>