

Mouse anti-Human. CD11c Monoclonal Antibody | anti-ITGAX antibody

CD11c

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62555	Mouse	Human.	Immunohistochemistry, Immunoprecipitation, Functional Assay, Immunofluorescence, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

CD11c; Monoclonal Antibody; CD11 antigen-like family member C; Complement component 3 receptor 4 subunit; Integrin alpha-X; integrin; alpha X (antigen CD11C (p150); alpha polypeptide); Leu M5 alpha subunit; Leukocyte adhesion glycoprotein p150 95 alpha chain; Myeloid membrane antigen alpha subunit; p150/95; anti-ITGAX antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human.
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	HC1/1
Specificity	CD11c, also known as Integrin, alpha X (complement component 3 receptor 4 subunit) (ITGAX), is a type I transmembrane protein found at high levels on most human dendritic cells, but also on monocytes, macrophages, neutrophils, and some B cells that induces cellular activation and helps trigger neutrophil respiratory burst; expressed in hairy cell leukemias, acute non-lymphocytic leukemias, and some B-cell chronic lymphocytic leukemias.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	1163
Applicable Applications for anti-ITGAX antibody	IHC (Immunohistochemistry), IP (Immunoprecipitation), FA (Functional Assay), IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ELISA
Localization	Cell surface
Positive Control	Human dendritic cells. Human lymph nodes and tonsils.
Human Chromosomal Location	16p11.2
Immunogen	Human monocytic U937 phorbol ester-treated cells
Quantity and Buffer	0.5 ml at 200ug/ml with BSA and azide
Preparation and Storage	Store at 2 to 8 degree C. Antibody is stable for 24 months.

ADDITIONAL INFORMATION

References	<ol style="list-style-type: none"> Nicolaou, F., et al. 2003. CD11c gene expression in hairy cell leukemia is dependent upon activation of the proto-oncogenes Ras and JunD. Blood 101: 4033-4041. Edwards, A.D., et al. 2003. Relationships among murine CD11c (high) dendritic cell subsets as revealed by baseline gene expression patterns. J. Immunol. 171: 47-60.
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	34452173
NCBI GeneID	3687
NCBI Accession #	NP_000878.2
NCBI GenBank Nucleotide #	NM_000887.4
UniProt Accession #	P20702; Q8IVA6
Molecular Weight	145kDa
NCBI Official Full Name	integrin alpha-X isoform 2
NCBI Official Synonym Full Names	integrin, alpha X (complement component 3 receptor 4 subunit)
NCBI Official Symbol	ITGAX
NCBI Official Synonym Symbols	CD11C; SLEB6
NCBI Protein Information	integrin alpha-X; leu M5, alpha subunit; p150 95 integrin alpha chain; CD11 antigen-like family member C; leukocyte adhesion receptor p150,95; myeloid membrane antigen, alpha subunit; leukocyte surface antigen p150,95, alpha subunit; leukocyte adhesion glycoprotein p150,95 alpha chain; integrin, alpha X (antigen CD11C (p150), alpha polypeptide)
UniProt Protein Name	Integrin alpha-X
UniProt Gene Name	ITGAX
UniProt Synonym Gene Names	CD11C
UniProt Entry Name	ITAX_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd11c-antibody-clone-hc1-1-immunohistochemistry-immunoprecipitation-functional-assay-immunofluorescence-flow-cytometry-elisa/62555>