

Mouse Bcl-2 Monoclonal Antibody | anti-BCL2 antibody

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62560	Mouse	Human.Does not react with Mouse and Rat	Immunohistochemistry, Western Blot, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Bcl-2; Monoclonal Antibody; Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Mouse Monoclonal Antibody; Apoptosis regulator Bcl-2; B-cell CLL/lymphoma-2; anti-BCL2 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Does not react with Mouse and Rat
Clonality	Monoclonal
Isotype	IgG
Clone Number	100/D5 + 124
Specificity	This antibody recognizes a protein of 25-26kDa, identified as the bcl-2 alpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 alpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of bcl-2 alpha protein, whereas the normal or hyperplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be used in distinguishing between those follicular lymphomas that express bcl-2 protein and the small number in which the neoplastic cells are bcl-2 negative.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	239
Applicable Applications for anti-BCL2 antibody	IHC (Immunohistochemistry), WB (Western Blot), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Outer mitochondrial membranes and endoplasmic reticulum as well as nuclear membranes.
Immunogen	A synthetic peptide, aa41-54 (GAAPAGIFSSQPG-Cys) of human Bcl-2 (100/D5 & 124)
Hu-Chromosome Location	18q21.33
Positive Control	Jurkat, K562, HL-60, or HeLa Cells. Tonsil or follicular lymphomas.
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.


IMAGES



WB (Western Blot)
Western Blot of Bcl-2 in human skin using Bcl-2 Monoclonal Antibody (100/D5 + 124).



Application Data
Intracellular staining of Bcl-2 in Jurkat cells using Bcl-2 Monoclonal Antibody (100/D5 + 124) (red) and isotype control (green).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human non-Hodgkin's Lymphoma stained with Bcl-2 Monoclonal Antibody (100/D5 + 124). Note nuclear membrane & cytoplasmic staining.

ADDITIONAL INFORMATION

References	Pezzella F et. al.. American Journal of Pathology, 1990, 137(2):225-32
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	72198189
NCBI GeneID	596
NCBI Accession #	NP_000624.2
NCBI GenBank Nucleotide #	NM_000633.2
UniProt Accession #	P10415; P10416; Q13842; Q16197; C9JHD5
Molecular Weight	25-26kDa
NCBI Official Full Name	apoptosis regulator Bcl-2 alpha isoform
NCBI Official Synonym Full Names	BCL2, apoptosis regulator
NCBI Official Symbol	BCL2
NCBI Official Synonym Symbols	Bcl-2; PPP1R50
NCBI Protein Information	apoptosis regulator Bcl-2
UniProt Protein Name	Apoptosis regulator Bcl-2

UniProt Gene Name	BCL2
UniProt Entry Name	BCL2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/bcl-2-antibody-clone-100-d5-124-immunohistochemistry-western-blot-immunofluorescence-flow-cytometry-elisa/62560>