

Mouse anti-Human CD34 Monoclonal Antibody | anti-CD34 antibody

CD34 (Hematopoietic Stem Cell & Endothelial Marker) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62677	Mouse	Human	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

CD34; Monoclonal Antibody; CD34 (Hematopoietic Stem Cell & Endothelial Marker) Mouse Monoclonal Antibody; Hematopoietic Progenitor Cell Antigen; HPCA1; Mucosialin; anti-CD34 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	HPCA1/763
Specificity	This MAb recognizes a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34. Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, CD34 is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. Its expression is also found in vascular endothelium. Additionally, it appears that proliferating endothelial cells express this molecule more than the non-proliferating endothelial cells. Anti-CD34 labels > 85% of angiosarcoma and KAP'si s sarcoma, but with a lower specificity.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.
Sequence Length	385
Applicable Applications for anti-CD34 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Cell Surface
Immunogen	Recombinant full-length human HPCA1 protein
Hu-Chromosome Location	1q32.2
Positive Control	KG-1 cells, Tonsil, or Angiosarcoma
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody is stable for 24 months.,Non-hazardous. No MSDS required.


IMAGES



IHC (Immunohistochemisry)
Formalin-fixed, paraffin-embedded human Placenta stained with CD34 Monoclonal Antibody (HPCA1/763)



IHC (Immunohiostchemistry)
Formalin-fixed, paraffin-embedded human Tonsil stained with CD34 Monoclonal Antibody (HPCA1/763)



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Angiosarcoma stained with CD34 Monoclonal Antibody (HPCA1/763)

ADDITIONAL INFORMATION

References	Sato T et al. Blood 94:2548-2554 (1999). Felshow DM et al. Blood 97:3768-3775 (2001).
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	68342038
NCBI GeneID	947
NCBI Accession #	NP_001020280.1
NCBI GenBank Nucleotide #	NM_001025109.1
UniProt Accession #	P28906; Q15970; Q15971; Q5JTA3; Q5JTA4; Q9UJB1; A8K664
Molecular Weight	90-110kDa
NCBI Official Full Name	hematopoietic progenitor cell antigen CD34 isoform a
NCBI Official Synonym Full Names	CD34 molecule
NCBI Official Symbol	CD34
NCBI Protein Information	hematopoietic progenitor cell antigen CD34
UniProt Protein Name	Hematopoietic progenitor cell antigen CD34
UniProt Gene Name	CD34
UniProt Entry Name	CD34_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd34-antibody-clone-hpca1-763-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/62677>