

Mouse CD86 Monoclonal Antibody | anti-CD86 antibody

CD86 (Dendritic Cells Maturation Marker) Mouse Monoclonal Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62712	Mouse	Human, Mouse, Rat	Immunohistochemistry, Western Blot, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS


CD86; Monoclonal Antibody; CD86 (Dendritic Cells Maturation Marker) Mouse Monoclonal Antibody; Activation B7-2 antigen; B lymphocyte activation antigen B72; CD28 antigen ligand 2; CD28LG2; CLS1; CTLA-4 counter-receptor B7.2; Early T-cell co-stimulatory molecule 1; ETC-1; FUN-1; LAB72; Ly-58; MB7-2; TS/A-2; anti-CD86 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	SPM600
Specificity	Recognizes a protein of 70kDa, which is identified as CD86. CD86 is a type I transmembrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is expressed at high levels on resting peripheral monocytes and dendritic cells and at very low density on resting B and T lymphocytes. CD86 expression is rapidly upregulated by B cell specific stimuli with peak expression at 18 to 42 hours after stimulation. CD86, along with CD80/B71, is an important accessory molecule in T cell co-stimulation via its interaction with CD28 and CD152/CTLA4. Since CD86 has rapid kinetics of induction, it is believed to be the major CD28 ligand expressed early in the immune response. It is also found on malignant Hodgkin and Reed Sternberg (HRS) cells in Hodgkin's disease.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide.
Sequence Length	217
Applicable Applications for anti-CD86 antibody	IHC (Immunohistochemistry), WB (Western Blot), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Cell Surface
Immunogen	Recombinant human full-length CD86 protein
Hu-Chromosome Location	3q13.33
Positive Control	293T or Jukat cells. Monocytes and dendritic cells, activated T, B and natural killer cells in lymph node or tonsil.
Preparation and Storage	Antibody with azide - store at 2°C to 8°C. Non-hazardous. No MSDS required.



IF (Immunofluorescence)
Immunofluorescence staining of Ramos cells using CD86 Mouse Monoclonal Antibody (SPM600).; followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Red dot.



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Melanoma stained with CD86 Monoclonal Antibody (SPM600).

NCBI AND UNIPROT INFORMATION

NCBI GI #	332634950
NCBI GeneID	942
NCBI Accession #	NP_001193853.1
NCBI GenBank Nucleotide #	NM_001206924.1
UniProt Accession #	P42081; Q13655; Q6FHB1; Q6GTS4; Q7M4L5; A0N0P0; B7Z2F3; B7Z702; E7ETN5; E9PC27
Molecular Weight	70kDa
NCBI Official Full Name	T-lymphocyte activation antigen CD86 isoform 4
NCBI Official Synonym Full Names	CD86 molecule
NCBI Official Symbol	CD86
NCBI Official Synonym Symbols	B70; B7-2; B7.2; LAB72; CD28LG2
NCBI Protein Information	T-lymphocyte activation antigen CD86
UniProt Protein Name	T-lymphocyte activation antigen CD86
UniProt Gene Name	CD86
UniProt Synonym Gene Names	CD28LG2
UniProt Entry Name	CD86_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd86-antibody-clone-spm600-immunohistochemistry-western-blot-immunofluorescence-flow-cytometry-elisa/62712>