

Mouse anti-Human Cytokeratin 7 (KRT7) Monoclonal Antibody | anti-KRT7 antibody

Cytokeratin 7 (KRT7) (Glandular and Transitional Epithelial Marker) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62799	Mouse	Human	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Cytokeratin 7 (KRT7); Monoclonal Antibody; Cytokeratin 7 (KRT7) (Glandular and Transitional Epithelial Marker) Mouse Monoclonal Antibody; CK-7; K2C7; Keratin 55K Type II Cytoskeletal; Keratin Simple Epithelial Type 1 K7; Keratin Type II Cytoskeletal 7; Krt2-7; KRT7; Sarcolectin; SCL; Type II Mesothelial Keratin K7; Type-II Keratin Kb7; anti-KRT7 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG
Clone Number	KRT7/760 + KRT7/903
Specificity	It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This MAb is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This MAb is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-).
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	469
Applicable Applications for anti-KRT7 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Cytoplasmic
Immunogen	Recombinant full-length human KRT7 protein
Hu-Chromosome Location	12q13.13
Positive Control	HeLa cells. Carcinoma of ovary, lung, cervix, or breast
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.


IMAGES



IHC (Immunohistochemisry)
Formalin-fixed, paraffin-embedded human Endometrial Carcinoma stained with Cytokeratin 7
Monoclonal Antibody (KRT7/760 + KRT7/903)



IHC (Immunohiostchemistry)
Formalin-fixed, paraffin-embedded human Lung Carcinoma stained with Cytokeratin 7
Monoclonal Antibody (KRT7/760 + KRT7/903)



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with Cytokeratin 7
Monoclonal Antibody (KRT7/760 + KRT7/903)

ADDITIONAL INFORMATION

References	Ramaekers F, van Niekerk C, Poels L, Schaafsma E, Huijsmans A, Robben H, et al. Use of monoclonal antibodies to keratin 7 in the differential diagnosis of adenocarcinomas. Am J Pathol 1990;136:641-5
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	67782365
NCBI GeneID	3855
NCBI Accession #	NP_005547.3
NCBI GenBank Nucleotide #	NM_005556.3
UniProt Accession #	P08729; Q92676; Q9BUD8; Q9Y3R7
Molecular Weight	55kDa
NCBI Official Full Name	keratin, type II cytoskeletal 7
NCBI Official Synonym Full Names	keratin 7
NCBI Official Symbol	KRT7
NCBI Official Synonym Symbols	K7; CK7; SCL; K2C7
NCBI Protein Information	keratin, type II cytoskeletal 7
UniProt Protein Name	Keratin, type II cytoskeletal 7
UniProt Gene Name	KRT7

UniProt Synonym Gene Names	SCL; CK-7; K7
UniProt Entry Name	K2C7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cytokeratin-7-krt7-antibody-clone-krt7-760-krt7-903-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/62799>