

Mouse anti-Human, Rat Cytokeratin 18 (KRT18) Monoclonal Antibody | anti-KRT18 antibody

Cytokeratin 18 (KRT18) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62808	Mouse	Human, Rat	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Cytokeratin 18 (KRT18); Monoclonal Antibody; Cytokeratin 18 (KRT18) Mouse Monoclonal Antibody; Cell Proliferation-inducing Gene 46 Protein; CK18; CYK18 Cytokeratin Endo B; K18; Keratin-18; Kerd; KRT18; anti-KRT18 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Rat
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	KRT18/1190
Specificity	This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs to the type A (acidic) subfamily of low molecular weight keratins, exists in combination with cytokeratin 8. It was reported that tissues from gastrointestinal tract are positive for both cytokeratin 8 and 18 but do not contain cytokeratin 14. Tissues from gastrointestinal tract, respiratory tract and urogenital tract, as well as endocrine and exocrine tissues and mesothelial cells are positive for cytokeratin 18.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	430
Applicable Applications for anti-KRT18 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Cytoplasmic
Immunogen	Recombinant human KRT18 protein
Hu-Chromosome Location	12q13.13
Positive Control	MCF-7, HeLa cells, Breast Cancer
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded Rat Colon stained with Cytokeratin 18 Monoclonal Antibody (KRT18/1190).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with Cytokeratin 18 Monoclonal Antibody (KRT18/1190).

ADDITIONAL INFORMATION

References	Lauerova et al. Hybridoma 7: 495-504 (1988). Kovaik et al. Int.J.Cancer Suppl.3: 50-5 (1988). Taylor -Papadimitriou et.al. J.Cell Sci. 94: 403-13 (1989). Vojtšek et al. Folia Biol. 35(6): 373-382 (1989).
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557888
NCBI GeneID	3875
NCBI Accession #	NP_000215.1
NCBI GenBank Nucleotide #	NM_000224.2
UniProt Accession #	P05783; Q53G38; Q5U0N8; Q9BW26
Molecular Weight	45kDa
NCBI Official Full Name	keratin, type I cytoskeletal 18
NCBI Official Synonym Full Names	keratin 18
NCBI Official Symbol	KRT18
NCBI Official Synonym Symbols	K18; CK-18; CYK18
NCBI Protein Information	keratin, type I cytoskeletal 18
UniProt Protein Name	Keratin, type I cytoskeletal 18
UniProt Gene Name	KRT18
UniProt Synonym Gene Names	CYK18; CK-18; K18
UniProt Entry Name	K1C18_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.