

Mouse Cytochrome C Monoclonal Antibody | anti-CYCS antibody

Cytochrome C (Mitochondrial Marker) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62814	Mouse	Human, Mouse, Rat, Horse, Dog, Pigeon, Frog, Drosophila.Others not known	Immunohistochemistry, Western Blot, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

Cytochrome C; Monoclonal Antibody; Cytochrome C (Mitochondrial Marker) Mouse Monoclonal Antibody; CYC; CYCS; HCS; THC4; anti-CYCS antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse, Rat, Horse, Dog, Pigeon, Frog, Drosophila. Others not known
Clonality	Monoclonal
Isotype	IgG2b, kappa
Clone Number	7H8.2C12
Specificity	It recognizes an epitope within amino acids 93-104 of pigeon Cytochrome C, a well-characterized mobile electron transport protein that is essential to energy conversion in all aerobic organisms. In mammalian cells, this highly conserved protein is normally localized to the mitochondrial inter-membrane space. More recent studies have identified cytosolic cytochrome c as a factor necessary for activation of apoptosis. During apoptosis, cytochrome c is trans-located from the mitochondrial membrane to the cytosol, where it is required for activation of caspase-3 (CPP32). Overexpression of Bcl-2 has been shown to prevent the translocation of cytochrome c, thereby blocking the apoptotic process. Overexpression of Bax has been shown to induce the release of cytochrome c and to induce cell death. The release of cytochrome c from the mitochondria is thought to trigger an apoptotic cascade, whereby Apaf-1 binds to Apaf-3 (caspase-9) in a cytochrome c-dependent manner, leading to caspase-9 cleavage of caspase-3. This MAbs recognizes total cytochrome C which includes both apocytochrome (i.e. cytochrome in the cytosol without heme attached) and holocytochrome (i.e cytochrome in the mitochondria with heme attached).
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 1mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	105
Applicable Applications for anti-CYCS antibody	IHC (Immunohistochemistry), WB (Western Blot), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Cytoplasmic
Immunogen	Synthetic peptides corresponding to amino acid 1-80, 81-104 and 66-104 of pigeon cytochrome c
Hu-Chromosome Location	7p15.3
Positive Control	K-562, HL-60, Jurkat, NIH3T3 or PC-3 cells. Liver or Cardiac muscle.
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohiostchemistry)
Formalin-fixed, paraffin-embedded human Pancreas stained with Cytochrome C Monoclonal Antibody (7H8.2C12).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Salivary Tumor stained with Cytochrome C Monoclonal Antibody (7H8.2C12).

ADDITIONAL INFORMATION

References	Goshorn SG, E Retzel, and R Jemmerson. Common Structural Features among Monoclonal Antibodies Binding the Same Antigenic Region of Cytochrome c. J Biol Chem 266:2134-2142 (1991).
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	11128019
NCBI GeneID	54205
NCBI Accession #	NP_061820.1
NCBI GenBank Nucleotide #	NM_018947.5
UniProt Accession #	P99999; P00001; Q6NUR2; Q6NX69; Q96BV4; A4D166; B2R4I1
Molecular Weight	15kDa
NCBI Official Full Name	cytochrome c
NCBI Official Synonym Full Names	cytochrome c, somatic
NCBI Official Symbol	CYCS
NCBI Official Synonym Symbols	CYC; HCS; THC4
NCBI Protein Information	cytochrome c
UniProt Protein Name	Cytochrome c
UniProt Gene Name	CYCS
UniProt Synonym Gene Names	CYC
UniProt Entry Name	CYC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.