

Mouse Ep-CAM /CD326 Monoclonal Antibody | anti-TACSTD1 antibody

Ep-CAM /CD326 (Epithelial Marker) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62825	Mouse	Human.Does not react with Rat	Immunohistochemistry, Western Blot, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Ep-CAM /CD326; Monoclonal Antibody; Ep-CAM /CD326 (Epithelial Marker) Mouse Monoclonal Antibody; Adenocarcinoma-associated Antigen; Cell Surface Glycoprotein Trop-1; EGP2; EGP314; EGP40; Epithelial Cell Adhesion Molecule; Epithelial Glycoprotein 314; ESA; KSA; TACD1; TROP1; Tumor-associated Calcium Signal Transducer 1 (TACSTD1); ECS-1; Epidermal Surface Antigen 1; ESA1; FLOT2; Flotillin-2; Membrane Component; Chromosome 17; Surface Marker-1 (M17S1); REG-1; Reggie-1; Reggie-2; anti-TACSTD1 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Does not react with Rat
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	MOC-31
Specificity	Binding epitope of this antibody is located in the first EGF-like repeat domain (EGF1) between amino acids 27-59 of Ep-CAM. EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	314
Applicable Applications for anti-TACSTD1 antibody	IHC (Immunohistochemistry), WB (Western Blot), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Cell Surface
Immunogen	Neuraminidase treated GLS-1 human small cell lung carcinoma cells
Hu-Chromosome Location	2q21
Positive Control	HT29 cells or breast tumor
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.


IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Endometrial Carcinoma stained with Ep-CAM Monoclonal Antibody (MOC-31).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with Ep-CAM Monoclonal Antibody (MOC-31).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with Ep-CAM Monoclonal Antibody (MOC-31).

ADDITIONAL INFORMATION

References	Myklebust AT, Beiske K, Pharo A, Davies CD, Aamdal S, Fodstad O: Selection of anti-SCLC antibodies for diagnosis of bone marrow metastasis. Br J Cancer Suppl 1991, 14:49-53 Manon J. Winter, Iris D. Nagtegaal, J. Han J. M. van Krieken, and Sergey V. Litvinov. The Epithelial Cell Adhesion Molecule (Ep-CAM) as a Morphoregulatory Molecule Is a Tool in Surgical Pathology. J Biol Chem 125:437-446 (1994)
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	218505670
NCBI GeneID	4072
NCBI Accession #	NP_002345.2
NCBI GenBank Nucleotide #	NM_002354.2
UniProt Accession #	P16422; P18180; Q6FG26; Q6FG49; Q96C47; Q9UCD0
Molecular Weight	40-43kDa
NCBI Official Full Name	epithelial cell adhesion molecule
NCBI Official Synonym Full Names	epithelial cell adhesion molecule
NCBI Official Symbol	EPCAM
NCBI Official Synonym Symbols	ESA; KSA; M4S1; MK-1; DIAR5; EGP-2; EGP40; KS1/4; MIC18; TROP1; EGP314; HNPCC8; TACSTD1
NCBI Protein Information	epithelial cell adhesion molecule

UniProt Protein Name	Epithelial cell adhesion molecule
UniProt Gene Name	EPCAM
UniProt Synonym Gene Names	GA733-2; M1S2; M4S1; MIC18; TACSTD1; TROP1; Ep-CAM; EGP; EGP314; hEGP314
UniProt Entry Name	EPCAM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/ep-cam-cd326-antibody-clone-moc-31-immunohistochemistry-western-blot-immunofluorescence-flow-cytometry-elisa/62825>