

Mouse p57Kip2 Monoclonal Antibody | anti-CDKN1C antibody

p57Kip2 (Mitotic Inhibitor/Suppressor Protein) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62827	Mouse	Human, Mouse. Others not known.	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

p57Kip2; Monoclonal Antibody; p57Kip2 (Mitotic Inhibitor/Suppressor Protein) Mouse Monoclonal Antibody; Beckwith Wiedemann syndrome (WBS); BWCR; Cyclin dependent kinase inhibitor 1C (CDKN1C); Cyclin dependent kinase inhibitor p57; KIP2; p57; anti-CDKN1C antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse. Others not known.
Clonality	Monoclonal
Isotype	IgG
Clone Number	KP10 + KIP2/880
Specificity	Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	316
Applicable Applications for anti-CDKN1C antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Nuclear
Immunogen	Recombinant human p57Kip2 protein
Hu-Chromosome Location	11p15.5
Positive Control	LS174T, Raji, HT29, SK-BR3 cells. Colon or Prostate carcinomas.
Preparation and Storage	Antibody with azide store at 2-8°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with p57 Monoclonal Antibody (KP10+KIP2/880).

ADDITIONAL INFORMATION

References	Lee, M.-H., et al. 1995. Cloning of p57, a cyclin-dependent kinase inhibitor with unique domain structure and tissue distribution. Genes Dev. 9: 639-649
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557441
NCBI GeneID	1028
NCBI Accession #	NP_000067.1
NCBI GenBank Nucleotide #	NM_000076.2
UniProt Accession #	P49918
Molecular Weight	57kDa
NCBI Official Full Name	cyclin-dependent kinase inhibitor 1C isoform a
NCBI Official Synonym Full Names	cyclin dependent kinase inhibitor 1C
NCBI Official Symbol	CDKN1C
NCBI Official Synonym Symbols	BWS; WBS; p57; BWCR; KIP2; p57Kip2
NCBI Protein Information	cyclin-dependent kinase inhibitor 1C
UniProt Protein Name	Cyclin-dependent kinase inhibitor 1C
UniProt Gene Name	CDKN1C
UniProt Synonym Gene Names	KIP2
UniProt Entry Name	CDN1C_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/p57kip2-antibody-clone-kp10-kip2-880-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/62827>