

Mouse NGF-Receptor (p75) /CD271 Monoclonal Antibody | anti-NGFR antibody

NGF-Receptor (p75) /CD271 (Soft Tissue Tumor Marker) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62852	Mouse	Human, Monkey, Baboon, Cat, Rabbit, Ferret. Does not react with Mouse and Rat. Others not known.	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay



PRODUCT SYNONYMS

NGF-Receptor (p75) /CD271; Monoclonal Antibody; NGF-Receptor (p75) /CD271 (Soft Tissue Tumor Marker) Mouse Monoclonal Antibody; CD271; Gp80-LNGFR; Low affinity nerve growth factor receptor; Low affinity neurotrophin receptor p75NTR; Nerve growth factor receptor (NGFR); TNFR superfamily member 16; p75 ICD; p75 Neurotrophin receptor; Tumor necrosis factor receptor superfamily member 16 (TNFRSF16); anti-NGFR antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human, Monkey, Baboon, Cat, Rabbit, Ferret. Does not react with Mouse and Rat. Others not known.
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	NGFR5
Specificity	It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). Its epitope spans in aa 1-160 of extracellular domain of NGFR/NTR. NGF-receptor contains an extracellular domain containing four 40-amino acid repeats with 6 cysteine residues at conserved positions followed by a serine/threonine-rich region, a single transmembrane domain, and a 155-amino acid cytoplasmic domain. The cysteine-rich region contains the nerve growth factor binding domain. NGF is important for the development, differentiation, and survival of variety of neuronal and non-neuronal cells. Its action is mediated by binding two distinct receptors, the high affinity p140 and low affinity p75.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	427
Applicable Applications for anti-NGFR antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Cell Surface and Cytoplasmic
Immunogen	NGFR from A875 melanoma cells
Hu-Chromosome Location	17q21-22
Positive Control	Neuronal axons, Schwann cells, and perineural cells of peripheral nerves, or tumors of nerve sheath differentiation e.g. Schwannoma, Neurofibroma. Soma and axons of sensory neurons, and ganglionic satellite cells. Melanomas.
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohiostchemistry)
Formalin-fixed, paraffin-embedded human Adrenal Gland stained with NGFR Mouse Monoclonal Antibody (NGFR5).

SDS-PAGE
SDS-PAGE Analysis Purified NGFR Mouse Monoclonal Antibody (NGFR5). Confirmation of Purity and Integrity of Antibody.



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Melanoma stained with NGFR Monoclonal Antibody (NGFR5).

ADDITIONAL INFORMATION

References	Marano N, et. al. Journal of Neurochemistry, 1987, 48:225-32
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505393
NCBI GeneID	4804
NCBI Accession #	NP_002498.1
NCBI GenBank Nucleotide #	NM_002507.3
UniProt Accession #	P08138; B2R961; B4E096
Molecular Weight	75kDa
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 16
NCBI Official Synonym Full Names	nerve growth factor receptor
NCBI Official Symbol	NGFR
NCBI Official Synonym Symbols	CD271; p75NTR; TNFRSF16; p75(NTR); Gp80-LNGFR
NCBI Protein Information	tumor necrosis factor receptor superfamily member 16
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 16

UniProt Gene Name	NGFR
UniProt Synonym Gene Names	TNFRSF16; NGF receptor
UniProt Entry Name	TNR16_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/ngf-receptor-p75-cd271-antibody-clone-ngfr5-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/62852>