

Mouse CD22 / BL-CAM Monoclonal Antibody | anti-CD22 antibody

CD22 / BL-CAM Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62889	Mouse	Human.Others not tested	Immunofluorescence, Flow Cytometry, Functional Assay, Functional Assay


PRODUCT SYNONYMS

CD22 / BL-CAM; Monoclonal Antibody; CD22 / BL-CAM Mouse Monoclonal Antibody; B-lymphocyte cell adhesion molecule (BL-CAM); B-cell receptor CD22; CD22; Lectin 2; Lyb8; Sialic acid-binding Ig-like lectin 2 (Siglec-2); SIGLEC2; T-cell surface antigen Leu-14; anti-CD22 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Others not tested
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	FR10B4
Specificity	Recognizes a protein of 130-140kDa, identified as CD22 (also known as BL-CAM). CD22 expression is restricted to normal and neoplastic B cells and is absent from other haemopoietic cell types. In B-cell ontogeny, CD22 is first expressed in the cytoplasm of pro-B and pre-B cells, and on the surface as B cells mature to become IgD+. It is not expressed by plasma cells, CD22 is found highly expressed in follicular mantle and marginal zone B-cells, and while germinal center B-cells are relatively weak. CD22 is a member of the immunoglobulin superfamily and serves as an adhesion receptor for sialic acid-bearing ligands expressed on erythrocytes and all leukocyte classes. It also associates with tyrosine kinases and play a role in signal transduction and B-cell activation.
Form/Format	1.0mg/ml of Ab purified from Bioreactor Concentrate by Protein A/G.
Sequence Length	759
Applicable Applications for anti-CD22 antibody	IF (Immunofluorescence), FCM/FACS (Flow Cytometry), FA (Functional Assay)
Cellular Localization	Cell Surface
Immunogen	Human Burkitt's lymphoma cells
Hu-Chromosome Location	19q13.1
Positive Control	Raji, Daudi, IM9, JY25 and human peripheral blood lymphocytes. Human tonsil and lymphocytes.
Preparation and Storage	Antibody without azide - store at -20 to -80°C. Non-hazardous. No MSDS required.

IMAGES



SDS-PAGE
SDS-PAGE Analysis Purified CD22 Mouse Monoclonal Antibody (FR10B4). Confirmation of Integrity and Purity of Antibody.

ADDITIONAL INFORMATION

References	Campana, D., et al., in : Knapp, W., et al. (eds), Leucocyte Typing IV, Oxford Univ. Press, pp 190-192
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	297374825
NCBI GeneID	933
NCBI Accession #	NP_001172028.1
NCBI GenBank Nucleotide #	NM_001185099.1
UniProt Accession #	P20273; O95699; O95701; O95702; O95703; Q01665; Q32M46; Q92872; Q92873; Q9UQA6; F5GYU4; F5H7U3
Molecular Weight	130-140kDa
NCBI Official Full Name	B-cell receptor CD22 isoform 2
NCBI Official Synonym Full Names	CD22 molecule
NCBI Official Symbol	CD22
NCBI Official Synonym Symbols	SIGLEC2; SIGLEC-2
NCBI Protein Information	B-cell receptor CD22
UniProt Protein Name	B-cell receptor CD22
UniProt Gene Name	CD22
UniProt Synonym Gene Names	SIGLEC2; BL-CAM; Siglec-2
UniProt Entry Name	CD22_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.