

Mouse CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) Monoclonal Antibody | anti-CFTR antibody

CFTR (Cystic Fibrosis Transmembrane Conductance Regulator)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62916	Mouse	Human. Others not known.	Immunohistochemistry, Western Blot, Immunofluorescence



PRODUCT SYNONYMS

CFTR (Cystic Fibrosis Transmembrane Conductance Regulator); Monoclonal Antibody; ABC35; ATP Binding Cassette Superfamily C Member 7 (ABCC7); cAMP-dependent chloride channel; CFTR; CFTR/MRP; Channel conductance-controlling ATPase; Cystic Fibrosis Transmembrane Conductance Regulator; MRP7; TNR CFTR; anti-CFTR antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Others not known.
Clonality	Monoclonal
Isotype	IgG1
Clone Number	CFTR/1341
Specificity	Recognizes a protein of 165-170kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structurally similar to multidrug resistance (Mdr1) protein and both are members of the superfamily of ATP-binding cassette (ABC) transporters, also known as traffic ATPases, which are implicated in the movement of various substrates. The CFTR protein is a small conductance adenosine 3', 5'-cyclic monophosphate (cAMP)-activated chloride ion channel found in the apical membranes of epithelia within the pancreas, airway, intestine, bile duct, sweat gland, and male genital ducts. CFTR is a valuable marker of human pancreatic duct cell development and differentiation.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	1480
Applicable Applications for anti-CFTR antibody	IHC (Immunohistochemistry), WB (Western Blot), IF (Immunofluorescence)
Immunogen	Recombinant human full-length CFTR
Cellular Localization	Cell Surface and Cytoplasmic
Hu-chromosome Location	7q31.2
Positive Control	MOLT-4 cells. Pancreas, Kidney or Placenta.
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Pancreas stained with CFTR Monoclonal Antibody (CFTR/1341).

SDS-PAGE
SDS-PAGE Analysis of Purified Bcl-6 Rabbit Recombinant Monoclonal Antibody (BCL6/1951R).

ADDITIONAL INFORMATION

References	Riordan, J.R., et al. 1989. Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. Science 245: 1066-1073.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	90421313
NCBI GeneID	1080
NCBI Accession #	NP_000483.3
NCBI GenBank Nucleotide #	NM_000492.3
UniProt Accession #	P13569; Q20BG8; Q20BH2; Q2I0A1; Q2I102
Molecular Weight	165-170kDa
NCBI Official Full Name	cystic fibrosis transmembrane conductance regulator
NCBI Official Synonym Full Names	cystic fibrosis transmembrane conductance regulator
NCBI Official Symbol	CFTR
NCBI Official Synonym Symbols	CF; MRP7; ABC35; ABCC7; CFTR/MRP; TNR-CFTR; dJ760C5.1
NCBI Protein Information	cystic fibrosis transmembrane conductance regulator
UniProt Protein Name	Cystic fibrosis transmembrane conductance regulator
UniProt Gene Name	CFTR
UniProt Synonym Gene Names	ABCC7; CFTR

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.