

Rabbit Chromogranin A/CHGA Polyclonal Antibody | anti-CHGA antibody

Chromogranin A/CHGA (Neuroendocrine Marker)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62931	Rabbit	Human.Does not react with rat. Others not known.	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Chromogranin A/CHGA; Polyclonal Antibody; Chromogranin A/CHGA (Neuroendocrine Marker); Beta-Granin; CGA; CHGA; Chromogranin A Parathyroid Secretory Protein 1; ER-37; Pancreastatin; Parastatin; Parathyroid Secretory Protein 1; Pituitary Secretory Protein I; SP-I; Vasostatin I or II; anti-CHGA antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human. Does not react with rat. Others not known.
Clonality	Polyclonal
Isotype	IgG, kappa
Specificity	Chromogranin A is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, pheochromocytomas, paragangliomas, and other neuroendocrine tumors. Co-expression of chromogranin A and neuron specific enolase (NSE) is common in neuroendocrine neoplasms. Reportedly, co-expression of certain keratins and chromogranin indicates neuroendocrine lineage. The presence of strong anti-chromogranin staining and absence of anti-keratin staining should raise the possibility of paraganglioma. The co-expression of chromogranin and NSE is typical of neuroendocrine neoplasms. Most pituitary adenomas and prolactinomas readily express chromogranin.
Form/Format	200ug/ml of Ab purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	457
Applicable Applications for anti-CHGA antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Immunogen	Recombinant human full-length Chromogranin A protein
Cellular Localization	Finely granular Cytoplasmic
Hu-chromosome Location	14q32.12
Positive Control	PC12 cells. Adrenal gland, bowel, parathyroid, pancreas, or pheochromocytoma.
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Pancreas stained with Chromogranin A Rabbit Polyclonal Antibody



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Parathyroid stained with Chromogranin A Rabbit Polyclonal Antibody

ADDITIONAL INFORMATION

References	Konecki DS et. al. J Biol Chem 1987; 262:17026-30. Lloyd RV et. al. Am J Pathol 1988; 130:296-304.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502805
NCBI GeneID	1113
NCBI Accession #	NP_001266.1
NCBI GenBank Nucleotide #	NM_001275.3
UniProt Accession #	P10645; Q53FA8; Q6NR84; Q96E84; Q96GL7; Q9BQB5; B2R9E9
Molecular Weight	68-75kDa
NCBI Official Full Name	chromogranin-A isoform 1 preproprotein
NCBI Official Synonym Full Names	chromogranin A
NCBI Official Symbol	CHGA
NCBI Official Synonym Symbols	CGA
NCBI Protein Information	chromogranin-A
UniProt Protein Name	Chromogranin-A
UniProt Gene Name	CHGA
UniProt Synonym Gene Names	CgA; SP-I

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/chromogranin-a-chga-antibody-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/62931>