

Mouse Catenin, beta (CTNNB1) Monoclonal Antibody | anti-CTNNB1 antibody

Catenin, beta (CTNNB1)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62943	Mouse	Human, Mouse, Rat, Cow, Dog, Chicken. Others not known.	Immunohistochemistry, Western Blot, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Catenin; beta (CTNNB1); Monoclonal Antibody; Cadherin associated protein; beta 1 88kDa; Catenin beta-1; CATNB; CHBCAT; CTNNB1; anti-CTNNB1 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse, Rat, Cow, Dog, Chicken. Others not known.
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	rCTNNB1/2173
Specificity	Beta-catenin associates with the cytoplasmic portion of E-cadherin, which is necessary for the function of E-cadherin as an adhesion molecule. In normal tissues, beta-catenin is localized to the membrane of epithelial cells, consistent with its role in the cell adhesion complex. In breast ductal neoplasia, beta-catenin is usually localized in cellular membranes. However, in lobular neoplasia, a marked redistribution of beta-catenin throughout the cytoplasm results in a diffuse cytoplasmic pattern. Immuno-staining of beta-catenin and E-cadherin is helps in the accurate identification of ductal and lobular neoplasms, including a distinction between low-grade ductal carcinoma in situ (DCIS) and lobular carcinoma. Additionally, some rectal and gastric adenocarcinomas demonstrate diffuse cytoplasmic beta-catenin staining and a lack of membranous staining, mimicking the staining pattern observed with lobular breast carcinomas.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	781
Applicable Applications for anti-CTNNB1 antibody	IHC (Immunohistochemistry), WB (Western Blot), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Immunogen	Recombinant full-length human CCNB1 protein
Cellular Localization	Cell Surface and Cytoplasmic
Hu-chromosome Location	3p22.1
Positive Control	HeLa or MCF-7 cells. Breast carcinoma
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.


IMAGES



WB (Western Blot)
Western blot analysis of Hela cell lysate using Beta-Catenin Mouse Recombinant Monoclonal Ab (rCTNNB1/2173).



SDS-PAGE
SDS-PAGE Analysis of Purified Beta-Catenin Mouse Recombinant Monoclonal Ab (rCTNNB1/2173).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Breast Cancer stained with Beta-Catenin Mouse Recombinant Monoclonal Antibody (rCTNNB1/2173).

ADDITIONAL INFORMATION

References	Mastracci TL et. al. Mod Path. 2005; 18:741-751. Sarrio D et. al. Oncogene. 2004; 23:3272-3283. Dabbs DJ et. al. Am J Surg Path. 2007; 31:427-437.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	148233338
NCBI GeneID	1499
NCBI Accession #	NP_001091679.1
NCBI GenBank Nucleotide #	NM_001098209.1
UniProt Accession #	P35222; Q8NEW9; Q8NI94; Q9H391; A8K1L7
Molecular Weight	92kDa
NCBI Official Full Name	catenin beta-1 isoform 1
NCBI Official Synonym Full Names	catenin beta 1
NCBI Official Symbol	CTNNB1
NCBI Official Synonym Symbols	EVR7; CTNNB; MRD19; armadillo
NCBI Protein Information	catenin beta-1
UniProt Protein Name	Catenin beta-1

UniProt Gene Name	CTNNB1
UniProt Synonym Gene Names	CTNNB

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/catenin-beta-ctnnb1-antibody-clone-rctnnb1-2173-immunohistochemistry-western-blot-immunofluorescence-flow-cytometry-elisa/62943>