

Mouse Glypican-3 (GPC3) Monoclonal Antibody | anti-GPC3 antibody

Glypican-3 (GPC3) (Hepatocellular Carcinoma Marker)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA62999	Mouse	Human, Rat. Others not known.	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

Glypican-3 (GPC3); Monoclonal Antibody; Glypican-3 (GPC3) (Hepatocellular Carcinoma Marker); DGSX; Glypican proteoglycan 3; GPC3; GTR2-2; Heparan sulphate proteoglycan; Intestinal protein OCI-5; MXR7; OCI-5; SDYS; Secreted glypican-3; SGBS1; anti-GPC3 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Rat. Others not known.
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	rGPC3/863
Specificity	Glypican-3 (GPC3) is a glycosylphosphatidyl inositol-anchored membrane protein, which may also be found in a secreted form. Anti-GPC3 has been identified as a useful tumor marker for the hepatocellular carcinoma (HCC), hepatoblastoma, melanoma, testicular germ cell tumors, and Wilms tumor. In patients with HCC, GPC3 is overexpressed in neoplastic liver tissue and elevated in serum, but is undetectable in normal liver, benign liver, and the serum of healthy donors. GPC3 expression is also found to be higher in HCC liver tissue than in cirrhotic liver or liver with focal lesions such as dysplastic nodules and areas of hepatic adenoma (HA) with malignant transformation. In the context of testicular germ cell tumors, GPC3 expression is up regulated in certain histologic subtypes, specifically yolk sac tumors and choriocarcinoma. A high level of GPC3 expression is also found in some types of embryonal tumors, such as Wilms tumor and hepatoblastoma, with a low or undetectable expression in normal adjacent tissue. In patients with thyroid cancer, expression of GPC3 is dramatically enhanced in certain types of cancers: 100% in follicular carcinoma and 70% in papillary carcinoma. Expression of GPC3 in follicular carcinoma is significantly higher than that of follicular adenoma. In contrast, GPC3 is not expressed in anaplastic carcinoma.
Form/Format	200ug/ml of recombinant MAb purified by Protein A/G. Prepared in 1mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	603
Applicable Applications for anti-GPC3 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Immunogen	Recombinant full-length human GPC3 protein
Cellular Localization	Cytoplasmic
Hu-chromosome Location	Xq26.2
Positive Control	293T cells or Hepatocellular carcinoma
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



SDS-PAGE
SDS-PAGE Analysis of Purified Glypican-3 Mouse Recombinant Monoclonal Antibody (rGPC3/863).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Hepatocellular Carcinoma stained with Glypican-3 Mouse Recombinant Monoclonal Antibody (rGPC3/863).

ADDITIONAL INFORMATION

References	Yan, B., et al. 2011. Expression and clinicopathologic significance of glypican 3 in hepatocellular carcinoma. Ann. Diagn. Pathol. 15: 162-169.
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	257471006
NCBI GeneID	2719
NCBI Accession #	NP_001158089.1
NCBI GenBank Nucleotide #	NM_001164617.1
UniProt Accession #	P51654; Q2L880; Q2L882; C9JLE3; G3V1R0
Molecular Weight	67kDa
NCBI Official Full Name	glypican-3 isoform 1
NCBI Official Synonym Full Names	glypican 3
NCBI Official Symbol	GPC3
NCBI Official Synonym Symbols	SGB; DGSX; MXR7; SDYS; SGBS; OCI-5; SGBS1; GTR2-2
NCBI Protein Information	glypican-3
UniProt Protein Name	Glypican-3
UniProt Gene Name	GPC3
UniProt Synonym Gene Names	OCI5

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.