

Human Endothelial Cell Adhesion Molecule (ESAM) ELISA Kit | ESAM elisa kit

Human Endothelial Cell Adhesion Molecule (ESAM) ELISA Kit

CATALOG #
AAA63308

REACTIVITY
Human


PRODUCT SYNONYMS

Endothelial Cell Adhesion Molecule (ESAM); Human Endothelial Cell Adhesion Molecule (ESAM) ELISA Kit; ESAM elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of ESAM. No significant cross-reactivity or interference between ESAM and analogues was observed.
Sequence Length	390
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.117ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ESAM were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ESAM were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ESAM elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of ESAM in human tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ESAM. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ESAM. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ESAM, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of ESAM in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	148664205
NCBI GeneID	90952
NCBI Accession #	NP_620411.2
NCBI GenBank Nucleotide #	NM_138961.2
UniProt Accession #	Q96AP7; Q96T50; B4DVN8
Molecular Weight	15,170 Da
NCBI Official Full Name	endothelial cell-selective adhesion molecule
NCBI Official Synonym Full Names	endothelial cell adhesion molecule
NCBI Official Symbol	ESAM
NCBI Official Synonym Symbols	W117m
NCBI Protein Information	endothelial cell-selective adhesion molecule
UniProt Protein Name	Endothelial cell-selective adhesion molecule
UniProt Gene Name	ESAM

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/endothelial-cell-adhesion-molecule-esam/63308>