

Mouse Cyclin Dependent Kinase Inhibitor 2A (CDKN2A) ELISA Kit | CDKN2A elisa kit

Mouse Cyclin Dependent Kinase Inhibitor 2A (CDKN2A) ELISA Kit

CATALOG #
AAA63422

REACTIVITY
Mouse


PRODUCT SYNONYMS

Cyclin Dependent Kinase Inhibitor 2A (CDKN2A); Mouse Cyclin Dependent Kinase Inhibitor 2A (CDKN2A) ELISA Kit; GIL; CDKN2A elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of CDKN2A. No significant cross-reactivity or interference between CDKN2A and analogues was observed.
Sequence Length	169
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.108ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CDKN2A were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CDKN2A were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CDKN2A elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CDKN2A in mouse tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CDKN2A. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CDKN2A. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CDKN2A, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CDKN2A in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	6753390
NCBI GeneID	12578
NCBI Accession #	NP_034007.1
NCBI GenBank Nucleotide #	NM_009877.2
UniProt Accession #	Q64364; Q4U255; Q9QXC7; Q9R051; P51480
Molecular Weight	17,941 Da
NCBI Official Full Name	cyclin-dependent kinase inhibitor 2A isoform 1
NCBI Official Synonym Full Names	cyclin-dependent kinase inhibitor 2A
NCBI Official Symbol	Cdkn2a
NCBI Official Synonym Symbols	Arf; p16; MTS1; Pctr1; p19ARF; p16INK4a; p19<ARF>; ARF-INK4a; INK4a-ARF; Ink4a/Arf; p16(INK4a)
NCBI Protein Information	cyclin-dependent kinase inhibitor 2A
UniProt Protein Name	Tumor suppressor ARF
UniProt Gene Name	Cdkn2a
UniProt Synonym Gene Names	ARF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/cyclin-dependent-kinase-inhibitor-2a-cdkn2a/63422>