

Mouse Cyclophilin A (CYPA) ELISA Kit | CYPA elisa kit

Mouse Cyclophilin A (CYPA) ELISA Kit

CATALOG #
AAA63465

REACTIVITY
Mouse


PRODUCT SYNONYMS

Cyclophilin A (CYPA); Mouse Cyclophilin A (CYPA) ELISA Kit; PPIA; CyP-A; CYPH; Peptidylprolyl Isomerase A; Peptidyl-Prolyl Isomerase A; Peptidyl-prolyl cis-trans isomerase A; Cyclosporin A-binding protein; CYPA elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of CYPA. No significant cross-reactivity or interference between CYPA and analogues was observed.
Sequence Length	164
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.065ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CYPA were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CYPA were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CYPA elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CYPA in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CYPA. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CYPA. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CYPA, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CYPA in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	6679439
NCBI GeneID	268373
NCBI Accession #	NP_032933.1
NCBI GenBank Nucleotide #	NM_008907.1
UniProt Accession #	P17742; Q9CWJ5; Q9R137
Molecular Weight	17,971 Da
NCBI Official Full Name	peptidyl-prolyl cis-trans isomerase A
NCBI Official Synonym Full Names	peptidylprolyl isomerase A
NCBI Official Symbol	Ppia
NCBI Official Synonym Symbols	Cphn; CypA; CyP-18; 2700098C05
NCBI Protein Information	peptidyl-prolyl cis-trans isomerase A
UniProt Protein Name	Peptidyl-prolyl cis-trans isomerase A
UniProt Gene Name	Ppia
UniProt Synonym Gene Names	PPIase A

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.