

Human Afamin (AFM) ELISA Kit | AFM elisa kit

Human Afamin (AFM) ELISA Kit

CATALOG #
AAA63546

REACTIVITY
Human


PRODUCT SYNONYMS

Afamin (AFM); Human Afamin (AFM) ELISA Kit; ALB2; ALBA; ALF; A-ALB; Alpha-Albumin; AFM elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of AFM. No significant cross-reactivity or interference between AFM and analogues was observed.
Sequence Length	599
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	12.5-800ng/mL
Sensitivity	< 5.92ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level AFM were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level AFM were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for AFM elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of AFM in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to AFM. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to AFM. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain AFM, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of AFM in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4501987
NCBI GeneID	173
NCBI Accession #	NP_001124.1
NCBI GenBank Nucleotide #	NM_001133.2
UniProt Accession #	P43652; Q32MR3; Q4W5C5; A8K3E1
Molecular Weight	69,069 Da
NCBI Official Full Name	afamin
NCBI Official Synonym Full Names	afamin
NCBI Official Symbol	AFM
NCBI Official Synonym Symbols	ALF; ALB2; ALBA
NCBI Protein Information	afamin
UniProt Protein Name	Afamin
UniProt Gene Name	AFM
UniProt Synonym Gene Names	ALB2; ALBA; Alpha-Alb

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.