

Rat 5'-Nucleotidase, Ecto (NT5E) ELISA Kit | NT5E elisa kit

Rat 5'-Nucleotidase, Ecto (NT5E) ELISA Kit

CATALOG #
AAA63616

REACTIVITY
Rat


PRODUCT SYNONYMS

5'-Nucleotidase; Ecto (NT5E); Rat 5'-Nucleotidase; Ecto (NT5E) ELISA Kit; SAC; HCA2; SACI; Sacy; NT5E elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of NT5E. No significant cross-reactivity or interference between NT5E and analogues was observed.
Sequence Length	576
Samples	Serum, plasma, tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.113ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level NT5E were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level NT5E were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NT5E elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of NT5E in rat serum, plasma, tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to NT5E. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to NT5E. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain NT5E, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of NT5E in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	148747417
NCBI GeneID	58813
NCBI Accession #	NP_067587.2
NCBI GenBank Nucleotide #	NM_021576.2
UniProt Accession #	Q66HL0; Q4G083; P21588
Molecular Weight	63,951 Da
NCBI Official Full Name	5'-nucleotidase
NCBI Official Synonym Full Names	5' nucleotidase, ecto
NCBI Official Symbol	Nt5e
NCBI Official Synonym Symbols	Nt5; CD73
NCBI Protein Information	5'-nucleotidase
UniProt Protein Name	5' nucleotidase, ecto
UniProt Gene Name	Nt5e

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.