

Mouse Reelin (RL) ELISA Kit | RL elisa kit

Mouse Reelin (RL) ELISA Kit

CATALOG #
AAA63686

REACTIVITY
Mouse


PRODUCT SYNONYMS

Reelin (RL); Mouse Reelin (RL) ELISA Kit; RELN; RL elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of RL. No significant cross-reactivity or interference between RL and analogues was observed.
Sequence Length	3459
Samples	Serum, plasma, tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	6.5pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level RL were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level RL were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for RL elisa kit	Intended Uses: This kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of RL in mouse serum, plasma, tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to RL. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to RL. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain RL, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of RL in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	882939033
NCBI GeneID	19699
NCBI Accession #	NP_001297393.1
NCBI GenBank Nucleotide #	NM_001310464.1
UniProt Accession #	Q60841; Q9CUA6; E9PZ78
Molecular Weight	383,200 Da
NCBI Official Full Name	reelin isoform 2
NCBI Official Synonym Full Names	reelin
NCBI Official Symbol	Reln
NCBI Official Synonym Symbols	rl; reeler
NCBI Protein Information	reelin
UniProt Protein Name	Reelin
UniProt Gene Name	Reln
UniProt Synonym Gene Names	RI

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/reelin-rl/63686>