

Human Junctional Adhesion Molecule 3 (JAM3) ELISA Kit | JAM3 elisa kit

Human Junctional Adhesion Molecule 3 (JAM3) ELISA Kit

CATALOG #
AAA63859

REACTIVITY
Human


PRODUCT SYNONYMS

Junctional Adhesion Molecule 3 (JAM3); Human Junctional Adhesion Molecule 3 (JAM3) ELISA Kit; JAM-C; JAMC; JAM3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of JAM3. No significant cross-reactivity or interference between JAM3 and analogues was observed.
Sequence Length	259
Samples	Serum, plasma and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	6.2pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level JAM3 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level JAM3 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for JAM3 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of JAM3 in human serum, plasma and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to JAM3. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to JAM3. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain JAM3, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of JAM3 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	328927057
NCBI GeneID	83700
NCBI Accession #	NP_001192258.1
NCBI GenBank Nucleotide #	NM_001205329.1
UniProt Accession #	Q9BX67; Q8WWL8; Q96FL1; B3KWG9
Molecular Weight	29,223 Da
NCBI Official Full Name	junctional adhesion molecule C isoform 2
NCBI Official Synonym Full Names	junctional adhesion molecule 3
NCBI Official Symbol	JAM3
NCBI Official Synonym Symbols	JAMC; JAM-2; JAM-3; JAM-C
NCBI Protein Information	junctional adhesion molecule C
UniProt Protein Name	Junctional adhesion molecule C
UniProt Gene Name	JAM3
UniProt Synonym Gene Names	JAM-C; JAM-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.