

Human B-Cell Leukemia/Lymphoma 2 (Bcl2) ELISA Kit | Bcl2 elisa kit

Human B-Cell Leukemia/Lymphoma 2 (Bcl2) ELISA Kit

CATALOG #
AAA63950

REACTIVITY
Human


PRODUCT SYNONYMS

B-Cell Leukemia/Lymphoma 2 (Bcl2); Human B-Cell Leukemia/Lymphoma 2 (Bcl2) ELISA Kit; APAF3; CASPASE9c; ICE-LAP6; MCH6; Apoptosis-Related Cysteine Peptidase;; Bcl2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Bcl2. No significant cross-reactivity or interference between Bcl2 and analogues was observed.
Sequence Length	239
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	1.56-100ng/mL
Sensitivity	< 0.57ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Bcl2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Bcl2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Bcl2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of Bcl2 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to Bcl2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to Bcl2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Bcl2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of Bcl2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	72198189
NCBI GeneID	596
NCBI Accession #	NP_000624.2
NCBI GenBank Nucleotide #	NM_000633.2
UniProt Accession #	P10415; P10416; Q13842; Q16197; C9JHD5
Molecular Weight	22,337 Da
NCBI Official Full Name	apoptosis regulator Bcl-2 alpha isoform
NCBI Official Synonym Full Names	BCL2, apoptosis regulator
NCBI Official Symbol	BCL2
NCBI Official Synonym Symbols	Bcl-2; PPP1R50
NCBI Protein Information	apoptosis regulator Bcl-2
UniProt Protein Name	Apoptosis regulator Bcl-2
UniProt Gene Name	BCL2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/b-cell-leukemia-lymphoma-2-bcl2/63950>